

Recombinant Human KAP1 protein (denatured) ab171679

1 图像

描述

产品名称	重组人KAP1蛋白(denatured)
纯度	> 90 % SDS-PAGE.
表达系统	Escherichia coli
Accession	Q13263
蛋白长度	Protein fragment
无动物成分	No
性质	Recombinant
种属	Human
序列	MGSSHHHHHSSGLVPRGSHMGSKLIYFQLHRALKMIVDPVE PHGEMKFQ WDLNAWTKSAEAFGKIVAERP GTNSTGPAPMAPP RAPGPLSK QGSGSSQP MEVQEGYGFSGDDPYSSAEPHVSGVKRSRSGEGEVSGLMRK VPRVSLER LDLDTADSQPPVFKVFPGSTTEDYNLIVIERGAAAAATGQP GTAPAGTP GAPPLAGMAIVKEEETEAAIGAPPTATEGPETKPVLMALAEG PGAEGPRL ASPSGSTSSGLEVVAPEGTSAPGGGPGTLDDSATICRVCQKP GDLVMCNQ CEFCFHL DCHLPALQDVPGE EWSCSLCHVLPDLKEEDGSLSL DGADSTGV VAKLSPANQRKCERVLLALFCHEPCRPLHQLATDSTFSLDQP GGTLDLTL IRARLQEKLSPPYSSPQEFAQDVGRMFKQFNKLTEDKADVQS IIGLQRFF ETRMNEAFGD
预测分子量	49 kDa including tags
氨基酸	366 to 802
标签	His tag N-Terminus

技术指标

Our **Abpromise guarantee** covers the use of **ab171679** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用 SDS-PAGE

形式 Liquid

制备和贮存

稳定性和存储 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

pH: 8.00

Constituents: 2.4% Urea, 0.32% Tris HCl, 10% Glycerol (glycerin, glycerine)

常规信息

功能 Nuclear corepressor for KRAB domain-containing zinc finger proteins (KRAB-ZFPs). Mediates gene silencing by recruiting CHD3, a subunit of the nucleosome remodeling and deacetylation (NuRD) complex, and SETDB1 (which specifically methylates histone H3 at 'Lys-9' (H3K9me)) to the promoter regions of KRAB target genes. Enhances transcriptional repression by coordinating the increase in H3K9me, the decrease in histone H3 'Lys-9 and 'Lys-14' acetylation (H3K9ac and H3K14ac, respectively) and the disposition of HP1 proteins to silence gene expression. Recruitment of SETDB1 induces heterochromatinization. May play a role as a coactivator for CEBPB and NR3C1 in the transcriptional activation of ORM1. Also corepressor for ERBB4. Inhibits E2F1 activity by stimulating E2F1-HDAC1 complex formation and inhibiting E2F1 acetylation. May serve as a partial backup to prevent E2F1-mediated apoptosis in the absence of RB1. Important regulator of CDKN1A/p21(CIP1). Has E3 SUMO-protein ligase activity toward itself via its PHD-type zinc finger.

组织特异性 Expressed in all tissues tested including spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes.

通路 Protein modification; protein sumoylation.

序列相似性 Belongs to the TRIM/RBCC family.
Contains 2 B box-type zinc fingers.
Contains 1 bromo domain.
Contains 1 PHD-type zinc finger.
Contains 1 RING-type zinc finger.

结构域 The HP1 box is both necessary and sufficient for HP1 binding.
The PHD-type zinc finger enhances CEBPB transcriptional activity. The PHD-type zinc finger, the HP1 box and the bromo domain, function together to assemble the machinery required for repression of KRAB domain-containing proteins. Acts as an intramolecular SUMO E3 ligase for autSUMOylation of bromodomain.
The RING-finger-B Box-coiled-coil/tripartite motif (RBCC/TRIM motif) is required for interaction with the KRAB domain of KRAB-zinc finger proteins. Binds four zinc ions per molecule. The RING finger and the N-terminal of the leucine zipper alpha helical coiled-coil region of RBCC are required for oligomerization.
Contains one Pro-Xaa-Val-Xaa-Leu (PxVxL) motif, which is required for interaction with chromoshadow domains. This motif requires additional residues -7, -6, +4 and +5 of the central Val which contact the chromoshadow domain.

翻译后修饰 Phosphorylated upon DNA damage, probably by ATM or ATR. ATM-induced phosphorylation on


Ser-824 represses sumoylation leading to the de-repression of expression of a subset of genes involved in cell cycle control and apoptosis in response to genotoxic stress. Dephosphorylation by the phosphatases, PPP1CA and PP1CB forms, allows sumoylation and expression of TRIM28 target genes.

Sumoylation/desumoylation events regulate TRIM28-mediated transcriptional repression. Sumoylation is required for interaction with CHD3 and SETDB1 and the corepressor activity. Represses and is repressed by Ser-824 phosphorylation. Enhances the TRIM28 corepressor activity, inhibiting transcriptional activity of a number of genes including GADD45A and CDKN1A/p21. Lys-554, Lys-779 and Lys-804 are the major sites of sumoylation. In response to Dox-induced DNA damage, enhanced phosphorylation on Ser-824 prevents sumoylation and allows de-repression of CDKN1A/p21.

细胞定位

Nucleus. Associated with centromeric heterochromatin during cell differentiation through CBX1.

图片



15% SDS-PAGE analysis of ab171679 (3 μ g).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors