

Recombinant human PARP3/IRT1 protein ab79638

2 图像

描述

产品名称	重组人PARP3/IRT1蛋白
生物活性	Specific Activity: 207.5 U/mg.
纯度	> 80 % SDS-PAGE. Affinity purified.
表达系统	Baculovirus infected Sf9 cells
蛋白长度	Full length protein
无动物成分	No
性质	Recombinant
种属	Human

技术指标

Our **Abpromise guarantee** covers the use of **ab79638** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Functional Studies SDS-PAGE
形式	Liquid
补充说明	Protein previously labeled as PARP3.

制备和贮存

稳定性和存储	Shipped on Dry Ice. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle. pH: 8.00 Constituents: 0.0462% (R*,R*)-1,4-Dimercaptobutan-2,3-diol, 0.395% Tris HCl, 0.05% Tween, 50% Glycerol (glycerin, glycerine), 0.58% Sodium chloride This product is an active protein and may elicit a biological response in vivo, handle with caution.
--------	--

常规信息

功能	Involved in the base excision repair (BER) pathway, by catalyzing the poly(ADP-ribosyl)ation of a
----	---

limited number of acceptor proteins involved in chromatin architecture and in DNA metabolism. This modification follows DNA damages and appears as an obligatory step in a detection/signaling pathway leading to the repair of DNA strand breaks. May link the DNA damage surveillance network to the mitotic fidelity checkpoint. Negatively influences the G1/S cell cycle progression without interfering with centrosome duplication. Binds DNA. May be involved in the regulation of PRC2 and PRC3 complex-dependent gene silencing.

组织特异性

Widely expressed; the highest levels are in the kidney, skeletal muscle, liver, heart and spleen; also detected in pancreas, lung, placenta, brain, leukocytes, colon, small intestine, ovary, testis, prostate and thymus.

序列相似性

Contains 1 PARP alpha-helical domain.
Contains 1 PARP catalytic domain.

结构域

According to PubMed:10329013, the N-terminal domain (54 amino acids) of isoform 2 is responsible for its centrosomal localization. The C-terminal region contains the catalytic domain.


翻译后修饰

Auto-poly(ADP)-ribosylation.

细胞定位


Nucleus. Cytoplasm > cytoskeleton > centrosome. Cytoplasm > cytoskeleton > centrosome > centriole. Core component of the centrosome. Preferentially localized to the daughter centriole throughout the cell cycle. According to PubMed:16924674, it is almost exclusively localized in the nucleus and appears in numerous small foci and a small number of larger foci whereas a centrosomal location has not been detected.

图片



PARP3/IRT1 assay - Specific activity 207.5 U/mg

Functional Studies - Recombinant human PARP3/IRT1 protein (ab79638)



ab79638 on SDS-PAGE, Molecular Weight 87 kDa

10% SDS-PAGE, Coomassie staining

Lane 1: ab79638 6µg

Lane 2: Protein marker

SDS-PAGE - Recombinant human PARP3/IRT1 protein (ab79638)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors