

Recombinant Mouse CD36 protein ab208474

1 图像

描述	
产品名称	重组小鼠CD36蛋白
纯度	> 95 % SDS-PAGE. Affinity purified
内毒素水平	< 1.000 Eu/μg
表达系统	Insect cells
Accession	<u>Q08857</u>
蛋白长度	Protein fragment
无动物成分	No
性质	Recombinant
种属	Mouse
序列	ADPGDMLIEKTIKREVVLEEGTTAFKNWVKTGTTVYRQFWIF DVQNPDDV AKNSSKIKVKQRPYTYRVRYLAKENITQDPEDHTVSFVQPN GAIFEPSL SVGTEDDNFTVLNLAVAAAAPHIYQNSFVQVVLNSLIKKSKSS MFQTRSLK ELLWGYKDPFLSLVPYPISTTVGVFYPPYNDTVDGVYKVFNGK DNISKVAI IESYKGKRNL SYWPSYCDMINGTDAASFPPFVEKSRTLRFSS SDICRSIY AVFGSEIDLKGIPVYRFVLPANAFASPLQNPDNHCFCTEKVI SNNCTSYG VLDIGKCKEGKPVYISLPHFLHASPDVSEPIEGLHPNEDEHR TYLDVEPI TGFTLQFAKRLQVNILVKPARKIEALKNLKRPYIVPILWLNE TGTIGDEK AEMFKTQVTGKIKHHHHHH
预测分子量	47 kDa including tags
氨基酸	30 to 439
标签	His tag C-Terminus
额外的序列信息	Extracellular domain. NP_031669
技术指标	

Our **Abpromise guarantee** covers the use of **ab208474** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用 SDS-PAGE

形式 Liquid

制备和贮存

稳定性和存储 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

pH: 7.40

Constituents: 10% Glycerol (glycerin, glycerine), 90% PBS


常规信息

功能 Multifunctional glycoprotein that acts as receptor for a broad range of ligands. Ligands can be of proteinaceous nature like thrombospondin, fibronectin, collagen or amyloid-beta as well as of lipidic nature such as oxidized low-density lipoprotein (oxLDL), anionic phospholipids, long-chain fatty acids and bacterial diacylated lipopeptides. They are generally multivalent and can therefore engage multiple receptors simultaneously, the resulting formation of CD36 clusters initiates signal transduction and internalization of receptor-ligand complexes. The dependency on coreceptor signaling is strongly ligand specific. Cellular responses to these ligands are involved in angiogenesis, inflammatory response, fatty acid metabolism, taste and dietary fat processing in the intestine (Probable). Binds long-chain fatty acids and facilitates their transport into cells, thus participating in muscle lipid utilization, adipose energy storage, and gut fat absorption (By similarity) (PubMed:18353783, PubMed:21610069). In the small intestine, plays a role in proximal absorption of dietary fatty acid and cholesterol for optimal chylomicron formation, possibly through the activation of MAPK1/3 (ERK1/2) signaling pathway (By similarity) (PubMed:18753675). Involved in oral fat perception and preferences (PubMed:22240721, PubMed:25822988). Detection into the tongue of long-chain fatty acids leads to a rapid and sustained rise in flux and protein content of pancreatobiliary secretions (By similarity). In taste receptor cells, mediates the induction of an increase in intracellular calcium levels by long-chain fatty acids, leading to the activation of the gustatory neurons in the nucleus of the solitary tract (By similarity). Important factor in both ventromedial hypothalamus neuronal sensing of long-chain fatty acid and the regulation of energy and glucose homeostasis (By similarity). Receptor for thrombospondins, THBS1 and THBS2, mediating their antiangiogenic effects (By similarity). As a coreceptor for TLR4:TLR6 heterodimer, promotes inflammation in monocytes/macrophages. Upon ligand binding, such as oxLDL or amyloid-beta 42, interacts with the heterodimer TLR4:TLR6, the complex is internalized and triggers inflammatory response, leading to NF-kappa-B-dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway, as well as IL1B secretion, through the priming and activation of the NLRP3 inflammasome (By similarity) (PubMed:20037584). Selective and nonredundant sensor of microbial diacylated lipopeptide that signal via TLR2:TLR6 heterodimer, this cluster triggers signaling from the cell surface, leading to the NF-kappa-B-dependent production of TNF, via MYD88 signaling pathway and subsequently is targeted to the Golgi in a lipid-raft dependent pathway (By similarity) (PubMed:16880211). (Microbial infection) Directly mediates cytoadherence of Plasmodium falciparum parasitized erythrocytes and the internalization of particles independently of TLR signaling.

疾病相关 Platelet glycoprotein IV deficiency

	Coronary heart disease 7
序列相似性	Belongs to the CD36 family.
翻译后修饰	N-glycosylated and O-glycosylated with a ratio of 2:1. Ubiquitinated at Lys-469 and Lys-472. Ubiquitination is induced by fatty acids such as oleic acid and leads to degradation by the proteasome (PubMed:21610069, PubMed:18353783). Ubiquitination and degradation are inhibited by insulin which blocks the effect of fatty acids (PubMed:18353783).
细胞定位	Cell membrane. Membrane raft. Golgi apparatus. Apical cell membrane. Upon ligand-binding, internalized through dynamin-dependent endocytosis.

图片



15% SDS-PAGE analysis of 3 µg of ab208474.

57-70 kDa (SDS-PAGE under reducing conditions).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors