

Biotinylated Goat Anti-Rabbit IgG (H+L) (Ready to use) ab64256

★★★★★ [1 Abreviews](#) [101 References](#) [1 图像](#)

概述

| | |
|-------|---|
| 产品名称 | Biotinylated山羊抗兔IgG (H+L) (Ready to use) |
| 经测试应用 | 适用于: IHC-P |
| 常规说明 | <p>Staining protocol</p> <p>Deparaffinize and rehydrate formalin-fixed paraffin-embedded tissue section.</p> <p>Add appropriate hydrogen peroxide blocking solution to cover the sections. Incubate for 10 minutes. Wash 2 times in buffer.</p> <p>Perform appropriate pretreatment if required. Wash 3 times in buffer.</p> <p>Apply appropriate protein blocking solution and incubate for 5 minutes at room temperature to block non specific background staining. Wash 1 time in buffer.</p> <p>Apply primary antibody and incubate according to manufacturer's protocol.</p> <p>Wash 4 times in buffer. Apply Ready to use Biotinylated Goat anti Rabbit IgG (ab64256) solution and incubate for 10-15 minutes at room temperature. Wash 4 times in buffer.</p> <p>Apply enzyme labelled streptavidin and incubate according to manufacturer's protocol.</p> <p>Rinse 4 times in buffer. Place slide in appropriate chromogenic substrate and incubate until desired reaction is achieved. Rinse 4 times in buffer.</p> <p>Add enough drops of Haematoxylin to cover the section. Incubate for 1 minute.</p> <p>Rinse 7-8 times in tap water. Add Mounting Medium to cover the section.</p> |

This reagent is intended to be used in a labeled streptavidin biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody which reacts with the primary antibody, enzyme labeled streptavidin, and substrate chromogen.

性能

| | |
|------|---|
| 形式 | Liquid |
| 存放说明 | Store at +4°C. |
| 存储溶液 | pH: 7.2 Preservative: 0.0985% Sodium azide Constituent: PBS |

| | |
|-----------|--|
| Reagent说明 | This reagent is intended to be used in a labeled streptavidin biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody |
|-----------|--|

which reacts with the primary antibody, enzyme labeled streptavidin, and substrate chromogen.

细胞定位

Secreted

细胞定位

Secreted

应用


The Abpromise guarantee

Abpromise™承诺保证使用ab64256于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论 | 说明 |
|-------|------|-------------------------------------|
| IHC-P | | Use at an assay dependent dilution. |

图片



Ab64256 was used with the primary antibody [ab3517](#) in IHC-P.

See Abreview on [ab3517](#).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Biotinylated Goat Anti-Rabbit IgG (H+L) (Ready to use) (ab64256)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors