

Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed ab96935

★★★★★ [1 Abreviews](#) [7 References](#) [3 图像](#)

概述

产品名称	驴抗山羊IgG H&L (DyLight® 488)预吸附二抗
宿主	Donkey
靶标种属	Goat
特异性	By immunoelectrophoresis and ELISA this antibody reacts specifically with Goat IgG and with light chains common to other Goat immunoglobulins. No antibody was detected against non immunoglobulin serum proteins. Reduced cross-reactivity to chicken, horse, human, mouse, pig, rabbit and rat was detected.
经测试应用	适用于: Flow Cyt, IHC-P, ICC/IF, WB
最小交叉反应	Chicken, Horse, Human, Mouse, Pig, Rabbit, Rat more details
偶联物	DyLight® 488. Ex: 493nm, Em: 518nm

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	pH: 6.8 Preservative: 0.09% Sodium azide Constituents: 0.2% BSA, PBS
纯度	Immunogen affinity purified
纯化说明	Antiserum was cross adsorbed using chicken, human, mouse, pig, rabbit and rat immunosorbents to remove cross reactive antibodies. This antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 488.
克隆	多克隆
同种型	IgG

应用


The Abpromise guarantee

Abpromise™ 承诺保证使用 ab96935 于以下的经测试应用

“应用说明”部分下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		1/50 - 1/200.
IHC-P		1/50 - 1/500.
ICC/IF		1/50 - 1/500.
WB		1/1000 - 1/2000. Predicted molecular weight: 36 kDa. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.

图片




All lanes : cocktail of mouse anti-hnRNP , rabbit anti-actin, and goat anti-GAPDH at 1 µg/ml

All lanes : HeLa

Secondary

All lanes : cocktail of DyLight® 680-conjugated donkey anti-mouse (green), DyLight® 800-conjugated donkey anti-rabbit (red), and DyLight® 488-conjugated donkey anti-goat (blue) (ab96935) at 0.5 µg/ml


Predicted band size: 36 kDa



Immunohistochemical analysis of Paraformaldehyde perfusion-fixed frozen human neuroblastoma xenograft into a murine host. The image is labelling LIMP2 with **ab87637** at 20°C for 1 hour at a dilution of 1/50 with 5% Donkey Serum. Permeabilization was done with 0.2% Triton. Secondary used was ab96935 at a 1/250 dilution.

Immunohistochemistry (PFA perfusion fixed frozen sections) - Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed (ab96935)

Image is courtesy of an AbReview submitted by Ms Françoise Geffroy.



Emission spectra of DyLight® fluorochromes available in our catalog.

Line colors represent the approximate visible colors of the wavelength maxima.

Donkey Anti-Goat IgG H&L (DyLight® 488) preadsorbed (ab96935)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors