

# Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed ab96919

★★★★★ [1 Abreviews](#) [20 References](#) [4 图像](#)

### 概述

产品名称	驴抗兔IgG H&L (DyLight® 488)预吸附二抗
宿主	Donkey
靶标种属	Rabbit
特异性	By immunoelectrophoresis and ELISA this antibody reacts specifically with Rabbit IgG and with light chains common to other Rabbit immunoglobulins. No antibody was detected against non immunoglobulin serum proteins. Reduced cross-reactivity to bovine, chicken, goat, horse, human, mouse, pig and rat was detected.
经测试应用	适用于: WB, IHC-P, ICC/IF, Flow Cyt
最小交叉反应	Chicken, Cow, Goat, Horse, Human, Mouse, Pig, Rat
偶联物	DyLight® 488. Ex: 493nm, Em: 518nm

[more details](#)

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	pH: 6.8 Preservative: 0.09% Sodium azide Constituents: 0.2% BSA, PBS
纯度	Immunogen affinity purified
纯化说明	Antiserum was cross adsorbed using bovine, chicken, goat, human, mouse, pig and rat immunosorbents to remove cross reactive antibodies. This antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 488.
克隆	多克隆
同种型	IgG


### 应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab96919于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000 - 1/20000. Predicted molecular weight: 36 kDa.
IHC-P		1/50 - 1/500.
ICC/IF	★★★★★ (1)	1/50 - 1/500.
Flow Cyt		1/50 - 1/200.

图片




Western blot - Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed (ab96919)

rabbit anti-actin antibody at 1 µg/ml + HeLa whole cell lysate at 30 µg

Secondary

Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed (ab96919) at 1/1000 dilution (Donkey anti-rabbit IgG heavy and light chain cross-adsorbed antibody)


**Predicted band size:** 36 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed (ab96919)

This image is courtesy of an abreview submitted by Carl Hobbs, Kings's College London, United Kingdom.


Glial fibrillary acidic protein (GFAP) staining in mouse Formalin Fixed Paraffin Embedded sections. For antigen retrieval HIER in citrate buffer was performed. Donkey polyclonal Secondary Antibody to Rabbit IgG - H&L (DyLight® 488), pre-adsorbed (ab96919) was used as a secondary antibody at a dilution of 1/50.



Immunohistochemistry (Frozen sections) - Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed (ab96919)

E14 mouse brain sample was stained with primary antibody (**ab66155**). The secondary antibody used was ab96919 (1:200 dilution, incubate for 1 hour at room temperature). The blue background was from DAPI.

*This image is courtesy of an anonymous Abreview*



Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed (ab96919)

Emission spectra of DyLight® fluorochromes available in our catalog.

Line colors represent the approximate visible colors of the wavelength maxima.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors