

## Product datasheet

# Rabbit Anti-Sheep IgG H&L (HRP) ab6747

★★★★★ [1 Abreviews](#) [55 References](#) [3 图像](#)

### 概述

产品名称	兔抗羊IgG H&L (HRP)
宿主	Rabbit
靶标种属	Sheep
经测试应用	<b>适用于:</b> IHC-P, WB, ELISA, Immunomicroscopy, Dot blot, ICC/IF, IHC-Fr
免疫原	Sheep IgG whole molecule
偶联物	HRP

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	Preservative: 0.01% Gentamicin sulphate Constituents: 0.88% Sodium chloride, 1% BSA, 0.42% Potassium phosphate
纯度	Affinity purified
纯化说明	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities.
共轭说明	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities
克隆	多克隆
同种型	IgG
常规说明	Horseradish Peroxidase (HRP).


### 应用

**The Abpromise guarantee** **Abpromise™**承诺保证使用ab6747于以下的经测试应用


“应用说明”部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use at an assay dependent concentration.
WB	★★★★★ (1)	1/1000 - 1/4000. PubMed: 15319485
ELISA		Use at an assay dependent dilution.
Immunomicroscopy		Use at an assay dependent concentration.
Dot blot		Use at an assay dependent concentration.
ICC/IF		1/500 - 1/2500.
IHC-Fr		Use at an assay dependent concentration.

图片




ab6747 was used at dilution 1/2000 with the primary antibody [ab7832](#) in WB. See the review on [ab7832](#).



Representative immunohistochemical staining of parietal peritoneum tissue sections displaying HA in brown color (A-D). HA is dramatically reduced in the peritoneum from mice treated with PDF (B) compared to tissue from untreated mice (A) and mice pretreated with H398 2 hours before application of PDF (C) or PBS (D) (Scale bar: 100µm). The secondary antibody was Rabbit Anti-Sheep IgG H&L (HRP) (ab6747) at 1/100 dilution.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Rabbit Anti-Sheep IgG H&L (HRP) (ab6747)

Kälble F et al. Selective Blocking of TNF Receptor 1 Attenuates Peritoneal Dialysis Fluid Induced Inflammation of the Peritoneum in Mice. *PLoS One* 11:e0163314 (2016).



Ab6747 was used at dilution 1/100 with the primary antibody [ab53842](#) in IHC-P. See the review on [ab53842](#).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Rabbit Anti-Sheep IgG H&L (HRP) (ab6747)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors