

HeLa-DFO treated (0.5mM, 24h) Nuclear Lysate ab180880

[1 References](#) [3 图像](#)


概述

产品名称	HeLa-DFO treated (0.5mM, 24h)核裂解物
常规说明	We recommend aliquoting the extracts into single use fractions and then storing them at -80°C. The cells were treated for 24 hours with a 0.5mM solution of Deferoxamine (DFO). DFO is used as a hypoxia-mimetic agent to stabilize Hypoxia Inducible Factor 1 (HIF-1). DFO stabilizes HIF-1 through the inhibition of Prolyl Hydroxylases (PHDs) which target HIF-1 through degradation. The mechanism of DFO inhibition is likely through the chelation of Fe ²⁺ bound to the active site of PHD which is required for enzymatic activity.

性能

Mycoplasma free	Yes
形式	Liquid
存放说明	Shipped on Dry Ice. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.
存储溶液	Constituent: 10% (R*,R*)-1,4-Dimercaptobutan-2,3-diol
裂解物说明	The cells were treated for 24 hours with a 0.5mM solution of Deferoxamine (DFO). DFO is used as a hypoxia-mimetic agent to stabilize Hypoxia Inducible Factor 1 (HIF-1). DFO stabilizes HIF-1 through the inhibition of Prolyl Hydroxylases (PHDs) which target HIF-1 through degradation. The mechanism of DFO inhibition is likely through the chelation of Fe ²⁺ bound to the active site of PHD which is required for enzymatic activity.
背景	HeLa cells are human epithelial cells from a fatal cervical carcinoma. The cell line was derived from cervical cancer cells taken from Henrietta Lacks, in 1951. Horizontal gene transfer from human papillomavirus 18 (HPV18) to human cervical cells created the HeLa genome which is different from either parent genome in various ways including its number of chromosomes. HeLa cells have a modal chromosome number of 82, with 4 copies of chromosome 12 and 3 copies of chromosomes 6, 8, and 17. HeLa cells are adherent cells (they stick to surfaces) and maintain contact inhibition in vitro.

图片




Western blot - HeLa-DFO treated (0.5mM, 24h)
Nuclear Lysate (ab180880)

All lanes : Simply Blue Coomassie Stain 20ml

Lane 1 : HeLa nuclear extract lysate (**ab150036**)

Lane 2 : HeLa-DFO treated (0.5mM, 24h) Nuclear Lysate
(ab180880)

Lysates/proteins at 10 µg per lane.



Western blot - HeLa-DFO treated (0.5mM, 24h)
Nuclear Lysate (ab180880)

All lanes : Anti-HIF-1 alpha antibody [H1alpha67] (**ab1**) at 5 µg/ml


Lane 1 : HeLa whole cell lysate (**ab150035**)

Lane 2 : HeLa-DFO treated (0.5mM, 24h) Nuclear Lysate
(ab180880)

Lysates/proteins at 40 µg per lane.

Secondary

All lanes : Goat Anti-Mouse IgG H&L (HRP) preadsorbed
(**ab97040**) at 1/10000 dilution



Western blot - HeLa-DFO treated (0.5mM, 24h)
Nuclear Lysate (ab180880)

All lanes : Anti-HIF-1 alpha antibody [EP1215Y] (**ab51608**) at 1/2000 dilution

Lane 1 : HeLa nuclear extract lysate (**ab150036**)

Lane 2 : HeLa-DFO treated (0.5mM, 24h) Nuclear Lysate (ab180880)

Lysates/proteins at 40 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Exposure time: 8 minutes

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors