

WI-38 nuclear extract lysate (forskolin stimulated) ab14912

1 图像


概述

产品名称	WI-38核extract裂解物(forskolin stimulated)
常规说明	The WI-38 nuclear extracts have been collected after a 30-minute incubation with or without forskolin (20 μM). The protein content has been determined by a Bradford-based assay.

性能

Mycoplasma free	Yes
形式	Liquid
存放说明	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.
存储溶液	pH: 7.50 Constituents: 0.0002% Sodium molybdate, 0.216% Beta glycerophosphate, 0.018% Sodium orthovanadate, 0.042% Sodium fluoride, 0.015% DTT, 0.48% HEPES, 0.003% EDTA, 20% Glycerol, 2.34% Sodium chloride With Protease inhibitor

图片



Each lot has been tested for CREB activation by using TransAM™ pCREB Kits. The signal intensity for CREB activation in each lot is compared to the signal intensity obtained with extracts from unstimulated WI-38 VA13 cells. After the signals are blanked, the ratio of the signals from stimulated cells over unstimulated cells must be above 3. This ratio may vary depending on the basal level of CREB activation in a given cell type.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors