

Product datasheet

Anti-CD45R antibody [351C5] ab20139

1 References

概述

产品名称	Anti-CD45R抗体[351C5]
描述	小鼠单克隆抗体[351C5] to CD45R
宿主	Mouse
特异性	Specific to CD45R. 351C5 can be used in typing of leukaemias and lymphomas and in research on functional subsets of B and T cells.
经测试应用	适用于: Flow Cyt, IP, IHC-Fr
种属反应性	与反应: Human
免疫原	Raji cells.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Constituent: PBS
纯度	Purified IgM
纯化说明	Purified by size exclusion.
克隆	单克隆
克隆编号	351C5
骨髓瘤	unknown
同种型	IgM
轻链类型	unknown

应用

Our [Abpromise guarantee](#) covers the use of **ab20139** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
Flow Cyt		Use at an assay dependent concentration. ab91545 - Mouse monoclonal IgM, is suitable for use as an isotype control with this antibody.
IP		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.

靶标

功能	Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN.
疾病相关	Defects in PTPRC are a cause of severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive (T(-)B(+)NK(+)) SCID [MIM:608971]. A form of severe combined immunodeficiency (SCID), a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity, leukopenia, and low or absent antibody levels. Patients present in infancy recurrent, persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development. Genetic variations in PTPRC are involved in multiple sclerosis susceptibility (MS) [MIM:126200]. MS is a neurodegenerative disorder characterized by the gradual accumulation of focal plaques of demyelination particularly in the periventricular areas of the brain. Peripheral nerves are not affected. Onset usually in third or fourth decade with intermittent progression over an extended period. The cause is still uncertain.
序列相似性	Belongs to the protein-tyrosine phosphatase family. Receptor class 1/6 subfamily. Contains 2 fibronectin type-III domains. Contains 2 tyrosine-protein phosphatase domains.
结构域	The first PTPase domain interacts with SKAP1.
翻译后修饰	Heavily N- and O-glycosylated.
细胞定位	Membrane. Membrane raft. Colocalized with DPP4 in membrane rafts.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors