

FITC Anti-Cytomegalovirus AD169 antibody ab30952

概述

产品名称	FITC荧光Anti-Cytomegalovirus AD169抗体
描述	FITC荧光山羊多克隆抗体to Cytomegalovirus AD169
宿主	Goat
偶联物	FITC. Ex: 493nm, Em: 528nm
特异性	ab30952 recognises Cytomegalovirus strain AD169.
经测试应用	适用于: ICC, IHC-P
种属反应性	与反应: Human cytomegalovirus
免疫原	Cytomegalovirus AD169
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	pH: 7.20 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
纯化说明	Chromatographically purified IgG fraction.
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab30952于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration.

靶标

相关性 Cytomegalovirus (CMV) is a member of the Herpes virus family. Members of this family have a characteristic virion structure. The double stranded DNA genome is contained within an icosahedral capsid which is embedded in a proteinaceous layer (tegument) and surrounded by a lipid envelope that is decorated with virus-specific glycoprotein spikes. The viral genes are coordinately expressed in groups at various times after infection. Early viral proteins are expressed in the nucleus of infected cells within 3 to 24 hours of infection prior to the commencement of viral DNA replication. This is followed by expression of the early intermediate genes, which encode enzymes required for viral DNA replication. After 48 to 72 hours, a number of late viral antigens may be demonstrated in the nuclei and cytoplasm of infected cells. Cytomegalovirus strain AD169 is a laboratory-adapted strain and appears to lack a 15kb region of the 200kb genome that is present in clinical isolates. This region contains 19 open reading frames whose functions have yet to be elucidated. AD169 is also unique in that it is unable to enter latency and nearly always assumes lytic growth upon infection.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors