

Product datasheet

Anti-DR5 antibody [MD5-1] (Phycoerythrin) ab95619

1 图像

概述

产品名称	Anti-DR5抗体[MD5-1] (Phycoerythrin)
描述	亚美尼亚仓鼠单克隆抗体[MD5-1] to DR5 (Phycoerythrin)
宿主	Armenian hamster
偶联物	Phycoerythrin. Ex: 488nm, Em: 575nm
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Mouse
免疫原	Mouse DR5.
阳性对照	Mouse DR5 transfected BHK cells.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	Preservative: 0.09% Sodium Azide Constituents: PBS, 150mM Sodium chloride, pH 7.2
纯度	Protein G purified
克隆	单克隆
克隆编号	MD5-1
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab95619** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
----	------	----

应用	Ab评论	说明
----	------	----

Flow Cyt Use 0.125-0.25µg for 10⁵⁻⁸ cells.
Staining the cell sample in a final volume of 100 µl is recommended.

[ab101026](#) - Armenian Hamster monoclonal IgG, is suitable for use as an isotype control with this antibody.

靶标

功能 Receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.


组织特异性 Widely expressed in adult and fetal tissues; very highly expressed in tumor cell lines such as HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus, stomach and throughout the intestinal tract; not detectable in brain.

疾病相关 Squamous cell carcinoma of the head and neck

序列相似性 Contains 1 death domain.
Contains 3 TNFR-Cys repeats.

细胞定位 Membrane.

图片



Flow cytometry staining of non-transfected (left) and mouse DR5 transfected (right) BHK cells with 0.125 µg of Golden Syrian Hamster IgG Isotype Control PE (open histogram) or 0.125 µg of ab95619 (filled histogram). Total viable cells were used for analysis.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors