

Product datasheet

Anti-EYA4 antibody ab93865

2 图像

概述

产品名称	Anti-EYA4抗体
描述	兔多克隆抗体to EYA4
宿主	Rabbit
经测试应用	适用于: WB, IP
种属反应性	与反应: Human 预测可用于: Rabbit, Guinea pig, Chimpanzee, Rhesus monkey, Gorilla, Orangutan
免疫原	Synthetic peptide from within residues: MEDSQLNEQ SVKKTCTESD VSQSQNSRSM EMQDLASPHT LVGGDTPGS, corresponding to N terminal amino acids 1-50 of Human EYA4 (NP_004091.3) Run BLAST with ExPASy Run BLAST with NCBI
阳性对照	HeLa whole cell lysate

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	Preservative: 0.09% Sodium Azide Constituents: Tris citrate/phosphate, pH 7-8
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用


Our [Abpromise guarantee](#) covers the use of **ab93865** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
WB		

应用	Ab评论	说明
IP		
应用说明	<p>IP: Use at 2-5µg/mg of lysate. WB: 1/1000 - 1/5000. Predicted molecular weight: 70 kDa.</p> <p>Not yet tested in other applications. Optimal dilutions/concentrations should be determined by the end user.</p>	
靶标		
功能	<p>Tyrosine phosphatase that specifically dephosphorylates 'Tyr-142' of histone H2AX (H2AXY142ph). 'Tyr-142' phosphorylation of histone H2AX plays a central role in DNA repair and acts as a mark that distinguishes between apoptotic and repair responses to genotoxic stress. Promotes efficient DNA repair by dephosphorylating H2AX, promoting the recruitment of DNA repair complexes containing MDC1. Its function as histone phosphatase probably explains its role in transcription regulation during organogenesis. May be involved in development of the eye.</p>	
组织特异性	<p>Highly expressed in heart and skeletal muscle.</p>	
疾病相关	<p>Defects in EYA4 are the cause of deafness autosomal dominant type 10 (DFNA10) [MIM:601316]. DFNA10 is a form of sensorineural hearing loss. Sensorineural deafness results from damage to the neural receptors of the inner ear, the nerve pathways to the brain, or the area of the brain that receives sound information.</p> <p>Defects in EYA4 are the cause of cardiomyopathy dilated type 1J (CMD1J) [MIM:605362]. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia. Patients are at risk of premature death.</p>	
序列相似性	<p>Belongs to the HAD-like hydrolase superfamily. EYA family.</p>	
细胞定位	<p>Cytoplasm. Nucleus.</p>	

图片



All lanes : Anti-EYA4 antibody (ab93865) at 0.4 µg/ml

Lane 1 : HeLa cell lysate at 50 µg


Lane 2 : HeLa cell lysate at 15 µg

Lane 3 : HeLa cell lysate at 5 µg

Developed using the ECL technique.

Predicted band size: 70 kDa

Exposure time: 3 minutes



3µg ab93865 were used to immunoprecipitate EYA4 from 1mg HeLa whole cell lysate. 20% IP was loaded and probed with 1µg/ml ab93865.

Detection: chemiluminescence with exposure time of 10 seconds.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors