

Product datasheet

Histone H3 (K9 methylation) Panel (H3, mono methyl K9, di methyl K9, tri methyl K9) ab113754

1 References 5 图像

概述

产品名称

Histone H3 (K9 methylation)组合(H3, mono methyl K9, di methyl K9, tri methyl K9)

产品概述

ab113754 is a Histone H3 (K9 methylation) Panel designed for the validation and characterization of the methylation state of Histone H3 on K9. Methylation of the canonical core histones can contribute to the formation of transcriptionally active and inactive chromatin in response to various signalling pathways and is a central modification for regulating epigenetic transitions in chromatin. Histone H3 methylation on K9 is a repressive modification, and H3 K9 tri-methylation has been shown to establish binding sites for heterochromatin protein 1.


性能

存放说明

Please refer to protocols.


组件	1 units
ab1220 - Anti-Histone H3 (di methyl K9) antibody [mAbcam 1220] - ChIP Grade	1 x 25µg
ab9045 - Anti-Histone H3 (mono methyl K9) antibody - ChIP Grade	1 x 25µg
ab8898 - Anti-Histone H3 (tri methyl K9) antibody - ChIP Grade	1 x 25µg
ab176842 - Anti-Histone H3 antibody - Nuclear Loading Control and ChIP Grade	1 x 10µl

图片




ChIP - Histone H3 (K9 methylation) Panel (H3, mono methyl K9, di methyl K9, tri methyl K9) (ab113754)

Chromatin was prepared from U2OS cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10 min. The ChIP was performed with 25 µg of chromatin, 2 µg of **ab1791** (blue), and 20 µl of protein A/G sepharose beads. No antibody was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Taqman approach). Primers and probes are located in the first kb of the transcribed region.




ChIP - Histone H3 (K9 methylation) Panel (H3, mono methyl K9, di methyl K9, tri methyl K9) (ab113754)

Chromatin was prepared from U2OS cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10 min. The ChIP was performed with 25 µg of chromatin, 2 µg of **ab9045** (blue), and 20 µl of protein A/G sepharose beads. No antibody was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Taqman approach for active and inactive loci, Sybr green approach for heterochromatic loci). Primers and probes are located in the first kb of the transcribed region.




Every new batch of this antibody is tested at Abcam in ChIP.

ChIP - Histone H3 (K9 methylation) Panel (H3, mono methyl K9, di methyl K9, tri methyl K9) (ab113754)



Every new batch of this antibody is tested at Abcam in ChIP.

ChIP - Histone H3 (K9 methylation) Panel (H3, mono methyl K9, di methyl K9, tri methyl K9) (ab113754)



Western blot - Histone H3 (K9 methylation) Panel (H3, mono methyl K9, di methyl K9, tri methyl K9) (ab113754)

Chromatin was prepared from U2OS cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10 min. The ChIP was performed with 25 µg of chromatin, 2 µg of [ab8898](#) (blue), and 20 µl of protein A/G sepharose beads. No antibody was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Taqman approach for active and inactive loci, Sybr green approach for heterochromatic loci). Primers and probes are located in the first kb of the transcribed region.

Chromatin was prepared from U2OS cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10min. The ChIP was performed with 25µg of chromatin, 2µg of [ab1220](#) (blue), and 20µl of Protein A/G sepharose beads. No antibody was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Taqman approach for active and inactive loci, Sybr green approach for heterochromatic loci). Primers and probes are located in the first Kb of the transcribed region.

All lanes : Anti-Histone H3 (mono methyl K9) antibody - ChIP Grade ([ab9045](#)) at 1 µg/ml

- Lane 2** : Human Histone H3 (unmodified) peptide ([ab7228](#))
- Lane 3** : Human Histone H3 (mono methyl K27) peptide ([ab1780](#))
- Lane 4** : Human Histone H3 (di methyl K27) peptide ([ab1781](#))
- Lane 5** : Human Histone H3 (tri methyl K27) peptide ([ab1782](#))
- Lane 6** : Human Histone H3 (mono methyl K4) peptide ([ab1340](#))
- Lane 7** : Human Histone H3 (mono methyl K9) peptide ([ab1771](#))

[ab9045](#) shows significantly greater reactivity with mono methyl K9. This can be seen in lane 7, as the addition of [ab1771](#) (mono methyl K9) completely blocks the activity of [ab9045](#). Weaker cross-reactivity is seen against mono methyl K27. This is shown in lane 3, as the addition of [ab1780](#) only partially blocks the activity of [ab9045](#).

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors