# abcam

## Product datasheet

## Anti-PADI2 / PAD2 antibody ab56075

## 1 图像

## 概述

产品名称 Anti-PADI2 / PAD2抗体

描述 兔多克隆抗体to PADI2 / PAD2

**宿主** Rabbit

经测试应用 适用于: WB, ELISA

种属反应性 与反应: Recombinant fragment

预测可用于: Mouse, Human ▲

免疫原 Synthetic peptide corresponding to C terminal residues of human PADI2/ PAD2 (peptidyl

arginine deiminase, type II)

性能

形式 Liquid

**存放说明** Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term.

存储溶液 Preservative: 0.01% Sodium Azide

Constituents: 50% Glycerol, PBS

纯**度** Immunogen affinity purified

纯**化**说明 Purified by antigen specific affinity chromatography.

 克隆
 多克隆

 同种型
 IgG

## 应用

Our Abpromise guarantee covers the use of ab56075 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| • • |    |    | • |  |  | • |  |
|-----|----|----|---|--|--|---|--|
| 应用  | Ab | 说明 |   |  |  |   |  |
|     | 评  |    |   |  |  |   |  |
|     |    |    |   |  |  |   |  |
|     | 论  |    |   |  |  |   |  |
|     |    |    |   |  |  |   |  |

## 应**用 Ab** 评

论

说明

WB

Use a concentration of 1  $\mu$ g/ml. Predicted molecular weight: 76 kDa. for 2 hours. This antibody has been tested in Western blot against the recombinant peptide used as an immunogen. We have no data on detection of endogenous protein.

**ELISA** 

Use at an assay dependent dilution.

#### 靶标

功能 Catalyzes the deimination of arginine residues of proteins.


序列相似性

Belongs to the protein arginine deiminase family.

细胞定位

Cytoplasm.

#### 图片



Western blot - Anti-PADI2 / PAD2 antibody (ab56075)

**All lanes :** Anti-PADI2 / PAD2 antibody (ab56075) at 1 μg/ml

Lane 1 : HeLa (Human epithelial carcinoma

cell line) Whole Cell Lysate

Lane 2: HepG2 (Human hepatocellular liver carcinoma cell line) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

### **Secondary**

**All lanes :** Goat polyclonal to Rabbit lgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

**Predicted band size:** 76 kDa **Observed band size:** 45 kDa

The length of PADI2 is currently uncertain.

According to Strausberg et al. (2002) [PMID: 12477932] it is 437aa in length (~45kDa).

However, according to Ishigami et al. (2002) [PMID: 12392711] PADI2 is 665aa (~75kDa).

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors