

Product datasheet

Recombinant Human BST2 protein ab109848

1 图像

概述

产品名称	重组人BST2蛋白
蛋白长度	Protein fragment

描述

性质	Recombinant
来源	Escherichia coli
氨基酸序列	
Accession	Q10589
种属	Human
序列	<p>MGSSHHHHHHSSGLVPRGSHMSEACRDGLRAVMECRN VTHLLQQLTEAQ KGFQDVEAQAATCNHTVMALMASLDAEKAQGQKKVEELEG EITTLNHKLQ DASAEVERLRRENQVLSVRIADKKYYPSSQDSS</p>
分子量	15 kDa including tags
氨基酸	50 to 161
标签	His tag N-Terminus

技术指标

Our [Abpromise guarantee](#) covers the use of **ab109848** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	SDS-PAGE Mass Spectrometry
质谱法	MALDI-TOF
纯度	> 80 % SDS-PAGE.
形式	Liquid

制备和贮存

稳定性和存储

Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

pH: 8.00

Constituents: 0.0154% DTT, 0.316% Tris HCl, 20% Glycerol, 0.058% Sodium chloride

常规信息

功能

May be involved in the sorting of secreted proteins (By similarity). May be involved in pre-B-cell growth. Antiretroviral defense protein, that blocks release of retrovirus from the cell surface. Depleted upon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) through ubiquitination and subsequent degradation. May play a role in B-cell activation in rheumatoid arthritis.

组织特异性

Predominantly expressed in liver, lung, heart and placenta. Lower levels in pancreas, kidney, skeletal muscle and brain. Overexpressed in multiple myeloma cells. Highly expressed during B-cell development, from pro-B precursors to plasma cells. Highly expressed on T-cells, monocytes, NK cells and dendritic cells (at protein level).

序列相似性

Belongs to the tetherin family.

结构域

The extracellular coiled coil domain is important for virus retention at the cell surface and prevention of virus spreading.


翻译后修饰

Monoubiquitinated by KSHV E3 ubiquitin-protein ligase K5, leading to its targeting to late endosomes and degradation.

细胞定位

Golgi apparatus > trans-Golgi network. Cell membrane. Cell membrane. Late endosome. Targeted to late endosomes upon KSHV infection and subsequent ubiquitination. Targeted to the trans-Golgi network by viral VPU protein.

图片



15% SDS-PAGE analysis of ab109848 (3 µg).

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors