

Product datasheet

SLBP peptide ab108380

概述

产品名称 SLBP多肽

描述

性质 Synthetic

氨基酸序列

序列 INYGKNTIAYDRIYIKEVPRHLRQPGIHPKTPNKFKKYSRRSVDQIQIKLWK

分子量 30 kDa

技术指标

Our [Abpromise guarantee](#) covers the use of **ab108380** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用 Blocking - Blocking peptide for Anti-SLBP antibody ([ab56027](#))

形式 Liquid

制备和贮存

稳定性和存储 Shipped at 4°C. Store at -20°C.

Preservative: None

Constituents: dH2O

常规信息

功能 RNA-binding protein involved in the histone pre-mRNA processing. Binds the stem-loop structure of replication-dependent histone pre-mRNAs and contributes to efficient 3'-end processing by stabilizing the complex between histone pre-mRNA and U7 small nuclear ribonucleoprotein (snRNP), via the histone downstream element (HDE). Plays an important role in targeting mature histone mRNA from the nucleus to the cytoplasm and to the translation machinery. Stabilizes mature histone mRNA and could be involved in cell-cycle regulation of histone gene expression. Involved in the mechanism by which growing oocytes accumulate histone proteins that support early embryogenesis. Binds to the 5' side of the stem-loop structure

	of histone pre-mRNAs.
组织特异性	Widely expressed.
序列相似性	Belongs to the SLBP family.
发展阶段	Regulated during the cell cycle: protein levels increase 10 to 20 fold in the late G1 and decrease at the S/G2 border.
结构域	Amino acids 31-34, 96-99 and 241-244 are necessary for interaction with the Importin alpha/Importin beta receptor. The first 18 amino acids, amino acids 69-76 and 179-182 are necessary for interaction with TNPO3. Amino acids 31-34, 96-99 and 241-244 are necessary for nuclear localization.
翻译后修饰	Phosphorylated on Thr-61 and Thr-62 in the S-phase. Phosphorylation of Thr-62 by CDK1 primes phosphorylation of Thr-61 by CK2. Phosphorylation of Thr-62 is required for its degradation by the proteasome at the end of the S phase. Its degradation is not required for histone mRNA degradation at the end of the S phase. All the phosphorylated forms detected are present in the cytoplasm. Both unphosphorylated and phosphorylated forms bind the stem-loop structure of histone mRNAs.
细胞定位	Cytoplasm. Nucleus. Polyribosome-associated. Localizes predominantly in the nucleus at the G1/G2 phases and the beginning of S phase. Through the S phase, partially redistributes to the cytoplasm. Binding to histone mRNA is necessary for cytoplasmic localization. Shuttles between the nucleus and the cytoplasm. Imported in the nucleus by the Importin alpha/Importin beta receptor.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors