

# Anti-SOX4 antibody ab80261

[35 References](#) [3 图像](#)

### 概述

产品名称	Anti-SOX4抗体
描述	兔多克隆抗体to SOX4
宿主	Rabbit
经测试应用	适用于: IHC-P, ICC/IF, WB
种属反应性	与反应: Human 预测可用于: Mouse, Rat, Rabbit, Goat, Horse, Chicken, Guinea pig, Cow, Dog, Drosophila melanogaster, Zebrafish 
免疫原	Synthetic peptide corresponding to Human SOX4 (N terminal). Immunogen located within the region from aa 38 and 87. Database link: <a href="#">Q06945</a>
阳性对照	WB: HeLa whole cell lysate ( <a href="#">ab150035</a> ). ICC/IF: SKNSH cells. IHC-P: Human testis tissue.

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	Preservative: 0.09% Sodium azide Constituents: 2% Sucrose, PBS
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

### 应用

**The Abpromise guarantee** **Abpromise™**承诺保证使用ab80261于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
IHC-P		Use a concentration of 5 µg/ml.

应用	Ab评论	说明
ICC/IF		Use a concentration of 1 µg/ml.
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 47 kDa (predicted molecular weight: 47 kDa). Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

## 靶标

<b>功能</b>	Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif.
<b>组织特异性</b>	Testis, brain, and heart.
<b>序列相似性</b>	Contains 1 HMG box DNA-binding domain.
<b>细胞定位</b>	Nucleus.

## 图片



Anti-SOX4 antibody (ab80261) at 1 µg/ml + HeLa cell lysate at 10 µg

### Secondary


HRP conjugated anti-Rabbit IgG at 1/50000 dilution

**Predicted band size:** 47 kDa

**Observed band size:** 47 kDa


**Additional bands at:** 75 kDa. We are unsure as to the identity of these extra bands.

Gel concentration 12%.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SOX4 antibody (ab80261)

ab80261 at 5 µg/ml staining SOX4 in human testis tissue by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections).



Immunocytochemistry/ Immunofluorescence - Anti-SOX4 antibody (ab80261)

ICC/IF image of ab80261 stained SKNSH cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab80261, 1µg/ml) overnight at +4°C. The secondary antibody (green) was [ab96899](#) Dylight 488 goat anti-rabbit IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors