

Product datasheet

Anti-SOX4 antibody ab80261

6 References 3 图像

概述

产品名称	Anti-SOX4抗体
描述	兔多克隆抗体to SOX4
宿主	Rabbit
经测试应用	适用于: IHC-P, ICC/IF, WB
种属反应性	与反应: Human 预测可用于: Mouse, Rat, Rabbit, Goat, Horse, Chicken, Guinea pig, Cow, Dog, Drosophila melanogaster, Zebrafish ▲
免疫原	Synthetic peptide corresponding to Human SOX4 (N terminal). Immunogen located within the region from aa 38 and 87. Database link: Q06945
阳性对照	WB: HeLa cell lysate. ICC/IF: SKNSH cells. IHC-P: Human testis tissue.

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	Preservative: 0.09% Sodium azide Constituents: 2% Sucrose, PBS
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab80261** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab 评论	说明
IHC-P		Use a concentration of 5 µg/ml.
ICC/IF		Use a concentration of 1 µg/ml.
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 47 kDa (predicted molecular weight: 47 kDa). Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

靶标

功能	Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif.
组织特异性	Testis, brain, and heart.
序列相似性	Contains 1 HMG box DNA-binding domain.
细胞定位	Nucleus.

图片



Anti-SOX4 antibody (ab80261) at 1 µg/ml + HeLa cell lysate at 10 µg

Secondary


HRP conjugated anti-Rabbit IgG at 1/50000 dilution

Predicted band size: 47 kDa

Observed band size: 47 kDa


Additional bands at: 75 kDa. We are unsure as to the identity of these extra bands.

Gel concentration 12%.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SOX4 antibody (ab80261)

ab80261 at 5 µg/ml staining SOX4 in human testis tissue by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections).



Immunocytochemistry/ Immunofluorescence - Anti-SOX4 antibody (ab80261)

ICC/IF image of ab80261 stained SKNSH cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab80261, 1µg/ml) overnight at +4°C. The secondary antibody (green) was [ab96899](#) Dylight 488 goat anti-rabbit IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors