

Product datasheet

Anti-STAM2 antibody ab63372

2 References 1 图像

概述

产品名称	Anti-STAM2抗体
描述	兔多克隆抗体to STAM2
宿主	Rabbit
经测试应用	适用于: ELISA, WB
种属反应性	与反应: Mouse, Human
免疫原	Synthetic non-phosphopeptide derived from human STAM2 around the phosphorylation site of tyrosine 192 (S-L-Y ^P -P-S).
阳性对照	NIH/3T3 cell extracts treated with EGF (200ng/ml, 30mins).

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	Preservative: 0.02% Sodium Azide Constituents: 50% Glycerol, PBS (without Mg ²⁺ and Ca ²⁺), 150mM Sodium chloride, pH 7.4
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用


Our [Abpromise guarantee](#) covers the use of **ab63372** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ELISA		1/5000.
WB		1/500 - 1/1000. Detects a band of approximately 58 kDa (predicted molecular weight: 58 kDa).

功能	Involves in intracellular signal transduction mediated by cytokines and growth factors. Upon IL-2 and GM-CSF stimulation, it plays a role in signaling leading to DNA synthesis and MYC induction. May also play a role in T-cell development. Involved in down-regulation of receptor tyrosine kinase via multivesicular body (MVBs) when complexed with HGS (ESCRT-0 complex). The ESCRT-0 complex binds ubiquitin and acts as sorting machinery that recognizes ubiquitinated receptors and transfers them to further sequential lysosomal sorting/trafficking processes.
组织特异性	Ubiquitously expressed.
序列相似性	Belongs to the STAM family. Contains 1 ITAM domain. Contains 1 SH3 domain. Contains 1 UIM (ubiquitin-interacting motif) repeat. Contains 1 VHS domain.
结构域	The VHS and UIM domains mediate the interaction with ubiquitinated proteins. The SH3 domain mediates the interaction with USP8. Contains one Pro-Xaa-Val-Xaa-Leu (PxVxL) motif, which is required for interaction with chromoshadow domains. This motif requires additional residues -7, -6, +4 and +5 of the central Val which contact the chromoshadow domain.
翻译后修饰	Phosphorylated in response to IL-2, GM-CSF, EGF and PDGF.
细胞定位	Cytoplasm. Early endosome membrane.

图片



All lanes : Anti-STAM2 antibody (ab63372) at 1/500 dilution

Lane 1 : NIH/3T3 cell extracts treated with EGF (200ng/ml, 30mins)

Lane 2 : NIH/3T3 cell extracts treated with EGF (200ng/ml, 30mins) with immunising peptide at 5 μ g

Lysates/proteins at 5 μ g per lane.

Predicted band size: 58 kDa

Observed band size: 58 kDa

Additional bands at: >117 kDa, 38 kDa. We are unsure as to the identity of these extra bands.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors