


Product datasheet

Anti-TPD52L2 antibody ab113952

4 图像

概述

<b>产品名称</b>	Anti-TPD52L2抗体
<b>描述</b>	山羊多克隆抗体to TPD52L2
<b>宿主</b>	Goat
<b>经测试应用</b>	<b>适用于:</b> WB, ELISA, IHC-P
<b>种属反应性</b>	<b>与反应:</b> Human <b>预测可用于:</b> Cynomolgus monkey, Orangutan 
<b>免疫原</b>	Synthetic peptide: SGDKPLSDPAP with a Cysteine residue linker, corresponding to C terminal amino acids 195-205 of Human TPD52L2 (NP_003279.2).  <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
<b>阳性对照</b>	Human brain cortex, spleen and tonsil tissues; Human breast cancer lysate.

性能

<b>形式</b>	Liquid
<b>存放说明</b>	Shipped at 4°C. Store at -20°C.
<b>存储溶液</b>	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 0.5% BSA, 99% Tris buffered saline
<b>纯度</b>	Protein G purified
<b>克隆</b>	多克隆
<b>同种型</b>	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab113952** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


应用	Ab评论	说明
WB		Use a concentration of 0.1 - 0.3 µg/ml. Predicted molecular weight: 22 kDa.
ELISA		1/16000.
IHC-P		Use a concentration of 3.75 µg/ml.

## 靶标

### 序列相似性


Belongs to the TPD52 family.

## 图片




ab113952, at 3.75 µg/ml, staining TPD52L2 in Formalin-Fixed, Paraffin-Embedded Human brain cortex tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TPD52L2 antibody (ab113952)




ab113952, at 3.75 µg/ml, staining TPD52L2 in Formalin-Fixed, Paraffin-Embedded Human spleen tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TPD52L2 antibody (ab113952)



ab113952, at 3.75 µg/ml, staining TPD52L2 in Formalin-Fixed, Paraffin-Embedded Human tonsil tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TPD52L2 antibody (ab113952)



Anti-TPD52L2 antibody (ab113952) at 0.1 µg/ml + Human breast cancer lysate (in RIPA buffer) at 35 µg

Developed using the ECL technique.

**Predicted band size:** 22 kDa

Western blot - Anti-TPD52L2 antibody (ab113952)

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors