

## Product datasheet

# Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade ab15584

★★★★☆ 3 Abreviews 5 References 8 图像

### 概述

产品名称	Anti-Histone H3 (mono methyl R2)抗体- ChIP Grade
描述	兔多克隆抗体to Histone H3 (mono methyl R2) - ChIP Grade
宿主	Rabbit
经测试应用	适用于: IP, ChIP/Chip, Dot blot, IHC-P, PepArr, WB, ChIP
种属反应性	与反应: Mouse, Cow, Human, Saccharomyces cerevisiae
免疫原	Synthetic peptide within Histone H3 aa 1-100 (mono methyl R2) conjugated to Keyhole Limpet Haemocyanin (KLH). The exact sequence is proprietary. (Peptide available as <a href="#">ab1775</a> )
阳性对照	WB: Calf thymus histone, C3H10T1/2 and 293 lysates. ICC/IF: HeLa cells.

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: PBS, 1% BSA
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

### 应用

Our [Abpromise guarantee](#) covers the use of **ab15584** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
IP		Use a concentration of 5 µg/ml.
ChIP/Chip		Use at an assay dependent concentration.
Dot blot		Use at an assay dependent concentration.
IHC-P		1/80. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
PepArr		Use a concentration of 0.2 - 2 µg/ml.
WB	★★★★☆	Use a concentration of 0.5 - 2 µg/ml. Detects a band of approximately 17 kDa (predicted molecular weight: 17 kDa).
ChIP		Use 2-4 µg for 25 µg of chromatin.

## 靶标


<b>功能</b>	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
<b>序列相似性</b>	Belongs to the histone H3 family.
<b>发展阶段</b>	Expressed during S phase, then expression strongly decreases as cell division slows down during the process of differentiation.
<b>翻译后修饰</b>	<p>Acetylation is generally linked to gene activation. Acetylation on Lys-10 (H3K9ac) impairs methylation at Arg-9 (H3R8me2s). Acetylation on Lys-19 (H3K18ac) and Lys-24 (H3K24ac) favors methylation at Arg-18 (H3R17me).</p> <p>Citrullination at Arg-9 (H3R8ci) and/or Arg-18 (H3R17ci) by PADI4 impairs methylation and represses transcription.</p> <p>Asymmetric dimethylation at Arg-18 (H3R17me2a) by CARM1 is linked to gene activation. Symmetric dimethylation at Arg-9 (H3R8me2s) by PRMT5 is linked to gene repression. Asymmetric dimethylation at Arg-3 (H3R2me2a) by PRMT6 is linked to gene repression and is mutually exclusive with H3 Lys-5 methylation (H3K4me2 and H3K4me3). H3R2me2a is present at the 3' of genes regardless of their transcription state and is enriched on inactive promoters, while it is absent on active promoters.</p> <p>Methylation at Lys-5 (H3K4me), Lys-37 (H3K36me) and Lys-80 (H3K79me) are linked to gene activation. Methylation at Lys-5 (H3K4me) facilitates subsequent acetylation of H3 and H4. Methylation at Lys-80 (H3K79me) is associated with DNA double-strand break (DSB) responses and is a specific target for TP53BP1. Methylation at Lys-10 (H3K9me) and Lys-28 (H3K27me) are linked to gene repression. Methylation at Lys-10 (H3K9me) is a specific target for HP1 proteins (CBX1, CBX3 and CBX5) and prevents subsequent phosphorylation at Ser-11 (H3S10ph) and acetylation of H3 and H4. Methylation at Lys-5 (H3K4me) and Lys-80 (H3K79me) require preliminary monoubiquitination of H2B at 'Lys-120'. Methylation at Lys-10 (H3K9me) and Lys-28 (H3K27me) are enriched in inactive X chromosome chromatin.</p> <p>Phosphorylated at Thr-4 (H3T3ph) by GSG2/haspin during prophase and dephosphorylated during anaphase. Phosphorylation at Ser-11 (H3S10ph) by AURKB is crucial for chromosome</p>

condensation and cell-cycle progression during mitosis and meiosis. In addition phosphorylation at Ser-11 (H3S10ph) by RPS6KA4 and RPS6KA5 is important during interphase because it enables the transcription of genes following external stimulation, like mitogens, stress, growth factors or UV irradiation and result in the activation of genes, such as c-fos and c-jun. Phosphorylation at Ser-11 (H3S10ph), which is linked to gene activation, prevents methylation at Lys-10 (H3K9me) but facilitates acetylation of H3 and H4. Phosphorylation at Ser-11 (H3S10ph) by AURKB mediates the dissociation of HP1 proteins (CBX1, CBX3 and CBX5) from heterochromatin. Phosphorylation at Ser-11 (H3S10ph) is also an essential regulatory mechanism for neoplastic cell transformation. Phosphorylated at Ser-29 (H3S28ph) by MLTK isoform 1, RPS6KA5 or AURKB during mitosis or upon ultraviolet B irradiation. Phosphorylation at Thr-7 (H3T6ph) by PRKCBB is a specific tag for epigenetic transcriptional activation that prevents demethylation of Lys-5 (H3K4me) by LSD1/KDM1A. At centromeres, specifically phosphorylated at Thr-12 (H3T11ph) from prophase to early anaphase, by DAPK3 and PKN1. Phosphorylation at Thr-12 (H3T11ph) by PKN1 is a specific tag for epigenetic transcriptional activation that promotes demethylation of Lys-10 (H3K9me) by KDM4C/JMJD2C. Phosphorylation at Tyr-42 (H3Y41ph) by JAK2 promotes exclusion of CBX5 (HP1 alpha) from chromatin. Monoubiquitinated by RAG1 in lymphoid cells, monoubiquitination is required for V(D)J recombination (By similarity). Ubiquitinated by the CUL4-DDB-RBX1 complex in response to ultraviolet irradiation. This may weaken the interaction between histones and DNA and facilitate DNA accessibility to repair proteins.

#### 细胞定位

Nucleus. Chromosome.


#### 图片



ICC/IF image of ab15584 stained mouse endothelial cells. The cells were fixed with methanol, permeabilized in 0.5% Triton and blocked with 5% BSA for 1 hour at 22°C. A diluted Cy3 conjugated goat anti-rabbit was used as a secondary antibody.


Immunocytochemistry/ Immunofluorescence - Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584)

This image is courtesy of an Abreview submitted by Marion Meixner



ChIP - Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584)


Chromatin was prepared from HeLa cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10min. The ChIP was performed with 25µg of chromatin, 2µg of ab15584 (blue), and 20µl of Protein A/G sepharose beads. No antibody was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Taqman approach). Primers and probes are located in the first kb of the transcribed region.



Peptide Array - Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584)

All batches of ab15584 are tested in Peptide Array against peptides to different Histone H3 modifications. Six dilutions of each peptide are printed on to the Peptide Array in triplicate and results are averaged before being plotted on to a graph. Results show strong binding to Histone H3 - mono methyl R2 peptide (ab1775), indicating that this antibody specifically recognises the Histone H3 - mono methyl R2 modification.

1. ab1775 – Histone H3 - mono methyl R2 peptide
2. ab38488 – Histone H3 - symmetric di methyl R2
3. ab2853 – Histone H3 - asymmetric di methyl R2
4. ab17566 – Histone H3 - non-modified
5. ab33971 – Histone H3 - mono methyl R8
6. ab13845 – Histone H3 - mono methyl R17
7. ab16935 – Histone H3 - asymmetric di methyl R17
8. ab32948 – Histone H3 - symm



Western blot - Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584)

Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584) at 1 µg/ml + Calf Thymus Histone Preparation Nuclear Lysate at 0.5 µg

**Secondary**

Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Performed under reducing conditions.


**Predicted band size:** 17 kDa

**Exposure time:** 3 minutes

**All lanes :** Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584) at 0.5 µg/ml


- Lane 1 :** Calf thymus histone at 0.4 µg
- Lane 2 :** C3H10T1/2 histones at 0.4 µg
- Lane 3 :** 293 histones at 0.4 µg
- Lane 4 :** Yeast extracts from 1 x 10<sup>6</sup> cells

**Predicted band size:** 17 kDa



Western blot - Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584)

ab15584 specifically recognises mono-methylated Histone H3 at R2 in calf thymus (lane1), mouse (lane2) and human (lane3) histone preparations as well as in yeast whole cell extract (lane4).




Immunoprecipitation - Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584)

Histone H3 was immunoprecipitated using 0.5mg HeLa whole cell extract, 5µg of Rabbit polyclonal to Histone H3 and 50µl of protein G magnetic beads (+). No antibody was added to the control (-). The antibody was incubated under agitation with Protein G beads for 10min, HeLa whole cell extract lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.

Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab15584.

Secondary: Clean-Blot IP Detection Reagent (HRP) at 1/500 dilution.

Band: 17kDa; Histone H3




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584)

Image courtesy of [Human Protein Atlas](#)

ab15584 staining histone H3 mono methyl R2 in human testis, showing a distinct and strong nuclear staining pattern in ductus semiferus and Leydig cells. Paraffin embedded human skin tissue was incubated with ab15584 (1/80 dilution) for 30 mins at room temperature. Antigen retrieval was performed by heat induction in citrate buffer pH 6.

ab15584 was tested in a tissue microarray (TMA) containing a wide range of normal and cancer tissues as well as a cell microarray consisting of a range of commonly used, well characterised human cell lines. Further results for this antibody can be found at [www.proteinatlas.org](http://www.proteinatlas.org)



Immunocytochemistry/ Immunofluorescence - Anti-Histone H3 (mono methyl R2) antibody - ChIP Grade (ab15584)

ICC/IF image of ab15584 stained HeLa cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab15584, 5µg/ml) overnight at +4°C. The secondary antibody (green) was ab96899, a goat anti-rabbit DyLight® 488 (IgG; H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors