

## Product datasheet

### Human PAR6 peptide ab49775

#### 概述

---

产品名称 人PAR6多肽

#### 描述

---

性质 Synthetic

#### 技术指标

---

Our [Abpromise guarantee](#) covers the use of **ab49775** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

形式 Liquid

补充说明

- First try to dissolve a small amount of peptide in either water or buffer. The more charged residues on a peptide, the more soluble it is in aqueous solutions.
- If the peptide doesn't dissolve try an organic solvent e.g. DMSO, then dilute using water or buffer.
- Consider that any solvent used must be compatible with your assay. If a peptide does not dissolve and you need to recover it, lyophilise to remove the solvent.
- Gentle warming and sonication can effectively aid peptide solubilisation. If the solution is cloudy or has gelled the peptide may be in suspension rather than solubilised.
- Peptides containing cysteine are easily oxidised, so should be prepared in solution just prior to use.

#### 制备和贮存

---

稳定性和存储 Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

Information available upon request.

#### 常规信息

---

功能 Adapter protein involved in asymmetrical cell division and cell polarization processes. Probably involved in the formation of epithelial tight junctions. Association with PARD3 may prevent the interaction of PARD3 with F11R/JAM1, thereby preventing tight junction assembly. The PARD6-PARD3 complex links GTP-bound Rho small GTPases to atypical protein kinase C proteins.

<b>组织特异性</b>	Expressed in pancreas, skeletal muscle, brain and heart. Weakly expressed in kidney and placenta.
<b>序列相似性</b>	Belongs to the PAR6 family. Contains 1 OPR domain. Contains 1 PDZ (DHR) domain. Contains 1 pseudo-CRIB domain.
<b>结构域</b>	The pseudo-CRIB domain together with the PDZ domain is required for the interaction with Rho small GTPases. The OPR domain mediates interactions with MAP2K5. The PDZ domain mediates the interaction with CRB3.
<b>细胞定位</b>	Cytoplasm. Cell membrane. Cell projection > ruffle. Cell junction > tight junction. Colocalizes with GTP-bound CDC42 or RAC1 at membrane ruffles and with PARD3 and PRKCI at epithelial tight junctions.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors