

Product datasheet

Anti-HAUSP / USP7 antibody - C-terminal ab71746

2 图像

概述

产品名称	Anti-HAUSP / USP7抗体- C-terminal
描述	兔多克隆抗体to HAUSP / USP7 - C-terminal
宿主	Rabbit
经测试应用	适用于: ELISA, IHC-P, WB
种属反应性	与反应: Human
免疫原	Synthetic peptide corresponding to Human HAUSP/ USP7 (C terminal) conjugated to keyhole limpet haemocyanin.
阳性对照	T47D cell lysate and human hepatocarcinoma tissue.

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: 0.09% Sodium Azide Constituents: PBS
纯度	Protein G purified
纯化说明	ab71746 is purified through a protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
克隆	多克隆
同种型	IgG

应用

Our [Abpromise guarantee](#) covers the use of **ab71746** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
ELISA		1/1000.

应用	Ab评论	说明
IHC-P		1/50 - 1/100.
WB		1/100 - 1/500. Detects a band of approximately 145 kDa (predicted molecular weight: 129 kDa).

靶标

功能

Hydrolase that deubiquitinates target proteins such as FOXO4, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN and DAXX (PubMed:11923872, PubMed:15053880, PubMed:16964248, PubMed:18716620, PubMed:25283148). Together with DAXX, prevents MDM2 self-ubiquitination and enhances the E3 ligase activity of MDM2 towards p53/TP53, thereby promoting p53/TP53 ubiquitination and proteasomal degradation. Deubiquitinates p53/TP53, preventing degradation of p53/TP53, and enhances p53/TP53-dependent transcription regulation, cell growth repression and apoptosis (PubMed:25283148). Deubiquitinates p53/TP53 and MDM2 and strongly stabilizes p53/TP53 even in the presence of excess MDM2, and also induces p53/TP53-dependent cell growth repression and apoptosis. Deubiquitination of FOXO4 in presence of hydrogen peroxide is not dependent on p53/TP53 and inhibits FOXO4-induced transcriptional activity. In association with DAXX, is involved in the deubiquitination and translocation of PTEN from the nucleus to the cytoplasm, both processes that are counteracted by PML. Involved in cell proliferation during early embryonic development. Involved in transcription-coupled nucleotide excision repair (TC-NER) in response to UV damage: recruited to DNA damage sites following interaction with KIAA1530/UJSSA and promotes deubiquitination of ERCC6, preventing UV-induced degradation of ERCC6. Contributes to the overall stabilization and trans-activation capability of the herpesvirus 1 trans-acting transcriptional protein ICP0/VMW110 during HSV-1 infection. Involved in maintenance of DNA methylation via its interaction with UHRF1 and DNMT1: acts by mediating deubiquitination of UHRF1 and DNMT1, preventing their degradation and promoting DNA methylation by DNMT1 (PubMed:21745816). Exhibits a preference towards 'Lys-48'-linked ubiquitin chains. Increases regulatory T-cells (Treg) suppressive capacity by deubiquitinating and stabilizing the transcription factor FOXP3 which is crucial for Treg cell function (PubMed:23973222).

组织特异性

Widely expressed. Overexpressed in prostate cancer.

序列相似性

Belongs to the peptidase C19 family.
Contains 1 MATH domain.
Contains 1 USP domain.

结构域

The C-terminus plays a role in its oligomerization.


翻译后修饰

Isoform 1: Phosphorylated. Isoform 1 is phosphorylated at positions Ser-18 and Ser-963.
Isoform 2: Not phosphorylated.
Isoform 1: Polyneddylated. Isoform 2: Not Polyneddylated.
Isoform 1 and isoform 2: Not sumoylated.
Isoform 1 and isoform 2: Polyubiquitinated by herpesvirus 1 trans-acting transcriptional protein ICP0/VMW110; leading to its subsequent proteasomal degradation. Isoform 1: Ubiquitinated at Lys-869.

细胞定位

Nucleus. Cytoplasm. Nucleus, PML body. Present in a minority of ND10 nuclear bodies. Association with ICP0/VMW110 at early times of infection leads to an increased proportion of USP7-containing ND10. Colocalizes with ATXN1 in the nucleus. Colocalized with DAXX in speckled structures. Colocalized with PML and PTEN in promyelocytic leukemia protein (PML) nuclear bodies.

图片




Western blot - Anti-HAUSP / USP7 antibody - C-terminal (ab71746)

Anti-HAUSP / USP7 antibody - C-terminal (ab71746) at 1/100 dilution + T47D cell lysate at 12.5 µg

Predicted band size: 129 kDa

Observed band size: 145 kDa

A lot of other bands can be seen. We do not know about the nature of these.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HAUSP / USP7 antibody - C-terminal (ab71746)

ab71746 at 1/50 dilution, staining HAUSP / USP7 in human hepatocarcinoma by Immunohistochemistry, Formalin-fixed, Paraffin-embedded tissue, followed by peroxidase-conjugated secondary antibody and AEC staining.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors