

## Product datasheet

# Anti-c-Myc (phospho T58) antibody ab28842

★★★★☆ 1 Abreviews 8 References 3 图像

### 概述

产品名称	Anti-c-Myc (phospho T58)抗体
描述	兔多克隆抗体to c-Myc (phospho T58)
宿主	Rabbit
经测试应用	适用于: IHC-P, IP, ELISA, ICC/IF, WB
种属反应性	与反应: Human 预测可用于: Mouse
免疫原	Synthetic peptide corresponding to c-Myc.
阳性对照	<div style="border: 1px solid #ccc; padding: 5px; margin-bottom: 5px;"> <a href="#">购买相配的WB阳性对照</a>  <a href="#">Recombinant Human c-Myc protein &gt;</a> </div> Ovarian cancer cell lysate, breast carcinoma tissue.

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: 50% Glycerol, 0.87% Sodium chloride
纯度	Without Mg <sup>2+</sup> and Ca <sup>2+</sup> Immunogen affinity purified
纯化说明	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.
克隆	多克隆
同种型	IgG

### 应用

Our [Abpromise guarantee](#) covers the use of **ab28842** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Ab评论	说明
IHC-P		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
ELISA		1/1000.
ICC/IF		Use a concentration of 1 µg/ml.
WB	★★★★☆	1/500 - 1/1000. Predicted molecular weight: 49 kDa.

## 靶标

### 功能

Participates in the regulation of gene transcription. Binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Seems to activate the transcription of growth-related genes.

### 疾病相关

Note=Overexpression of MYC is implicated in the etiology of a variety of hematopoietic tumors. Note=A chromosomal aberration involving MYC may be a cause of a form of B-cell chronic lymphocytic leukemia. Translocation t(8;12)(q24;q22) with BTG1. Defects in MYC are a cause of Burkitt lymphoma (BL) [MIM:113970]. A form of undifferentiated malignant lymphoma commonly manifested as a large osteolytic lesion in the jaw or as an abdominal mass. Note=Chromosomal aberrations involving MYC are usually found in Burkitt lymphoma. Translocations t(8;14), t(8;22) or t(2;8) which juxtapose MYC to one of the heavy or light chain immunoglobulin gene loci.

### 序列相似性

Contains 1 basic helix-loop-helix (bHLH) domain.


### 翻译后修饰

Phosphorylated by PRKDC. Phosphorylation at Thr-58 and Ser-62 by GSK3 is required for ubiquitination and degradation by the proteasome. Phosphorylation at Ser-329 by PIM2 leads to the stabilization of MYC (By similarity). Phosphorylation at Ser-62 by CDK2 prevents Ras-induced senescence. Ubiquitinated by the SCF(FBXW7) complex when phosphorylated at Thr-58 and Ser-62, leading to its degradation by the proteasome. In the nucleoplasm, ubiquitination is counteracted by USP28, which interacts with isoform 1 of FBXW7 (FBW7alpha), leading to its deubiquitination and preventing degradation. In the nucleolus, however, ubiquitination is not counteracted by USP28, due to the lack of interaction between isoform 4 of FBXW7 (FBW7gamma) and USP28, explaining the selective MYC degradation in the nucleolus. Also polyubiquitinated by the DCX(TRUSS) complex.

### 细胞定位

Nucleus > nucleoplasm. Nucleus > nucleolus.

## 图片



**Lane 1 :** Anti-c-Myc (phospho T58) antibody (ab28842) at 1/500 dilution

**Lane 2 :** Anti-c-Myc (phospho T58) antibody (ab28842) at 1/500 dilution ( preincubated with synthesized non-phosphopeptide )

**Lane 3 :** Anti-c-Myc (phospho T58) antibody (ab28842) at 1/500 dilution ( preincubated with synthesized phosphopeptide )


**Lanes 1 & 3 :** human ovarian cancer cell lysate

**Lane 2 :** human ovarian cancer cell lysate

**Predicted band size:** 49 kDa


**Observed band size:** 51 kDa

**Additional bands at:** 120 kDa, 55 kDa, 86 kDa. We are unsure as to the identity of these extra bands.



ICC/IF image of [ab28864](#) stained HeLa cells.

The cells were 4% PFA fixed (10 min) and then incubated in 1% BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ([ab28864](#), 1µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Immunohistochemical analysis of paraffin-embedded breast carcinoma. Left: Using Myc (Phospho-Thr58) Antibody; Right: The same antibody preincubated with synthesized phosphopeptide.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-c-Myc (phospho T58) antibody (ab28842)

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors