

Anti-Collagen V antibody ab7046

★★★★★ [11 Abreviews](#) [50 References](#) [3 图像](#)

概述

产品名称	Anti-Collagen V抗体
描述	兔多克隆抗体to Collagen V
宿主	Rabbit
特异性	Negligible cross-reactivity with Type I, II, III, IV or VI collagens. Non-specific cross reaction of anti-collagen antibodies with other human serum proteins or non-collagen extracellular matrix proteins is negligible.
经测试应用	适用于: WB, Sandwich ELISA, IHC-P, IP, IHC-Fr, ICC
种属反应性	与反应: Mouse, Cow, Human
免疫原	Full length native protein (purified) corresponding to Collagen V aa 1-1745.
常规说明	<p>Some class specific anti-collagens may be specific for three-dimensional epitopes which may result in diminished reactivity with denatured collagen or formalin-fixed, paraffin embedded tissues.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.01% Sodium azide Constituents: 4.8% Sodium borate, 0.15% EDTA, 0.44% Sodium chloride
纯度	Immunogen affinity purified
纯化说明	Immunoaffinity chromatography using immobilized antigens followed by extensive cross-adsorption against other collagens, human serum proteins and non-collagen extracellular matrix proteins to remove any unwanted specificities.

克隆 多克隆
同种型 IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab7046于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/5000 - 1/10000. Detects a band of approximately 180 kDa (predicted molecular weight: 180 kDa). Native conditions are recommended.
Sandwich ELISA		Use at an assay dependent concentration. Can be used with a suitable biotinylated detection antibody such as Rabbit polyclonal to Collagen V (Biotin) (ab6582).
IHC-P	★★★★★ (6)	1/50 - 1/200.
IP		Use at an assay dependent concentration.
IHC-Fr	★★★★☆ (2)	Use at an assay dependent concentration.
ICC		Use at an assay dependent concentration. PubMed: 18385800

靶标

功能 Type V collagen is a member of group I collagen (fibrillar forming collagen). It is a minor connective tissue component of nearly ubiquitous distribution. Type V collagen binds to DNA, heparan sulfate, thrombospondin, heparin, and insulin.

疾病相关 Ehlers-Danlos syndrome 1
Ehlers-Danlos syndrome 2


序列相似性 Belongs to the fibrillar collagen family.
Contains 1 fibrillar collagen NC1 domain.
Contains 1 laminin G-like domain.

结构域 The C-terminal propeptide, also known as COLFI domain, have crucial roles in tissue growth and repair by controlling both the intracellular assembly of procollagen molecules and the extracellular assembly of collagen fibrils. It binds a calcium ion which is essential for its function.

翻译后修饰 Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.
Sulfated on 40% of tyrosines.

细胞定位 Secreted > extracellular space > extracellular matrix.

图片



Western blot - Anti-Collagen V antibody (ab7046)

Anti-Collagen V antibody (ab7046) at 1 µg/ml + Human pancreas tissue lysate - total protein (**ab29816**) at 10 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (**ab97080**) at 1/5000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 180 kDa

Observed band size: 180 kDa

Additional bands at: 122 kDa, 54 kDa. We are unsure as to the identity of these extra bands.


Exposure time: 8 minutes



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen V antibody (ab7046)

This image is courtesy of an anonymous Abreview.

ab7046 staining Collagen V in Human pancreas tissue by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded tissue sections). The sections were fixed in formaldehyde and subjected to heat-mediated antigen retrieval in citrate pH 6 prior to blocking with 0.25% casein for 5 minutes at 25°C. The primary antibody was diluted 1/75 in Tris-HCL and incubated with the sample for 30 minutes at 25°C. An HRP polymer-conjugated goat anti-rabbit antibody was used as the secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen V antibody (ab7046)

Immunohistochemical analysis of formalin-fixed paraffin-embedded human lung (left) and placenta (right) sections labelling collagen V with ab7046 at a dilution of 1/200 for 45 minutes at room temperature. An antigen retrieval step was performed with 0.01 M sodium citrate buffer pH 6.0 at 100°C for 20 mins.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors