

Anti-FOXP3 antibody [236A/E7] ab20034

重组

★★★★☆ [37 Abreviews](#) [537 References](#) [14 图像](#)

概述

产品名称	Anti-FOXP3抗体[236A/E7]
描述	小鼠单克隆抗体[236A/E7] to FOXP3
宿主	Mouse
特异性	The epitope recognized by FOXP3 antibody [236A/E7] (ab20034) is between amino acids 105-236. FOXP3 antibody [236A/E7] (ab20034) is expected to detect full length FOXP3 as well as both cleaved forms.
经测试应用	适用于: mIHC, IHC-P, WB
种属反应性	与反应: Human
免疫原	Recombinant full length protein corresponding to Human FOXP3. Database link: Q9BZS1
阳性对照	WB: Human mammary gland lysate, HEK-293 transfected with FOXP3 expression vector containing a GFP-Myc-tag, whole cell lysate. IHC-P: Human tonsil and thymus tissue. mIHC: Human breast cancer tissue.
常规说明	This product has switched from a hybridoma to recombinant production method on 20 th January 2021.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)
纯度	Protein A purified
克隆	单克隆
克隆编号	236A/E7
骨髓瘤	P3-NS1/1-Ag4-1
同种型	IgG1

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab20034于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
mlHC		Use at an assay dependent concentration.
IHC-P	★★★★★ (26)	1/500. Perform heat mediated antigen retrieval via the microwave method before commencing with IHC staining protocol.
WB	★★★★★ (1)	1/1000. Detects a band of approximately 50 kDa (predicted molecular weight: 47 kDa).

靶标

功能

Probable transcription factor. Plays a critical role in the control of immune response.

疾病相关

Defects in FOXP3 are the cause of immunodeficiency polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) [MIM:304790]; also known as X-linked autoimmunity-immunodeficiency syndrome. IPEX is characterized by neonatal onset insulin-dependent diabetes mellitus, infections, secretory diarrhea, trombocytopenia, anemia and eczema. It is usually lethal in infancy.


序列相似性

Contains 1 C2H2-type zinc finger.
Contains 1 fork-head DNA-binding domain.

细胞定位

Nucleus.

图片



Multiplex immunohistochemistry - Anti-FOXP3 antibody [236A/E7] (ab20034)

This image is courtesy of ImmunoAtlas.

Fluorescence multiplex immunohistochemical analysis of Human breast cancer tissue (formalin-fixed paraffin-embedded section).

Merged staining of Anti-PD-L1 ([ab251611](#); cyan; Opal™ 520), Anti-Granzyme B ([ab219803](#); yellow; Opal™ 540), Anti-PD1 ([ab251613](#); magenta; Opal™ 570), Anti-pan Cytokeratin ([ab264485](#); red; Opal™ 620), Anti-EpCAM ([ab225894](#); red; Opal™ 620), Anti-CD8 alpha ([ab251596](#); green; Opal™ 650) and Anti-FOXP3 ([ab96048](#); orange; Opal™ 690). EpCAM and pan-cytokeratin share the same dye and color. Dyes are pseudo-colored for better contrast of the markers.

The immunostaining was performed on a Leica Biosystems BOND® MAX instrument with an Opal™ 6-Plex Detection Kit (NEL821001KT, Akoya Biosciences®).

The section was incubated in six rounds of staining; sequentially for

ab251611 (1/750 dilution), **ab219803** (1/250 dilution), **ab251613** (1/750 dilution), **ab264485** (0.5 µg/ml), **ab225894** (1/1250 dilution), **ab251596** (1/1500 dilution) and **ab96048** (10 µg/ml); each using a separate fluorescent tyramide signal amplification system. EDTA based antigen retrieval (Leica Biosystems BOND® Epitope Retrieval Solution 2, pH 9.0, 20 minutes) was used in between rounds of tyramide signal amplification to remove the antibody from the previous round, to avoid any cross-reactivity. DAPI (dark blue) was used as a nuclear counter stain.

Microscopy and pseudocoloring of individual Opal™ dyes was performed using a Vectra 3 Imaging System (Akoya Biosciences®).


This data is courtesy of ImmunoAtlas and it can be found [here](#).

Anti-FOXP3 antibody [236A/E7] (ab20034) at 1/1000 dilution + Human mammary gland lysate at 20 µg

Predicted band size: 47 kDa



Western blot - Anti-FOXP3 antibody [236A/E7] (ab20034)



Multiplex immunohistochemistry - Anti-FOXP3 antibody [236A/E7] (ab20034)

This image is courtesy of ImmunoAtlas.

Fluorescence multiplex immunohistochemical analysis of Human breast cancer tissue (formalin-fixed paraffin-embedded section).

Merged staining of Anti-PD-L1 (**ab251611**; cyan; Opal™ 520), Anti-Granzyme B (**ab219803**; yellow; Opal™ 540), Anti-PD1 (**ab251613**; magenta; Opal™ 570), Anti-pan Cytokeratin (**ab264485**; red; Opal™ 620), Anti-EpCAM (**ab225894**; red; Opal™ 620), Anti-CD8 alpha (**ab251596**; green; Opal™ 650) and Anti-FOXP3 (**ab96048**; orange; Opal™ 690). EpCAM and pan-cytokeratin share the same dye and color. Dyes are pseudo-colored for better contrast of the markers.


The immunostaining was performed on a Leica Biosystems BOND® MAX instrument with an Opal™ 6-Plex Detection Kit (NEL821001KT, Akoya Biosciences®).

The section was incubated in six rounds of staining; sequentially for **ab251611** (1/750 dilution), **ab219803** (1/250 dilution), **ab251613**

(1/750 dilution), **ab264485** (0.5 µg/ml), **ab225894** (1/1250 dilution), **ab251596** (1/1500 dilution) and **ab96048** (10 µg/ml); each using a separate fluorescent tyramide signal amplification system. EDTA based antigen retrieval (Leica Biosystems BOND[®] Epitope Retrieval Solution 2, pH 9.0, 20 minutes) was used in between rounds of tyramide signal amplification to remove the antibody from the previous round, to avoid any cross-reactivity. DAPI (dark blue) was used as a nuclear counter stain.

Microscopy and pseudocoloring of individual Opal[™] dyes was performed using a Vectra 3 Imaging System (Akoya Biosciences[®]).

This data is courtesy of ImmunoAtlas and it can be found [here](#).



Multiplex immunohistochemistry - Anti-FOXP3 antibody [236A/E7] (ab20034)

This image is courtesy of ImmunoAtlas.

Fluorescence multiplex immunohistochemical analysis of Human breast cancer tissue (formalin-fixed paraffin-embedded section).


Merged staining of Anti-PD-L1 (**ab251611**; cyan; Opal[™] 520), Anti-Granzyme B (**ab219803**; yellow; Opal[™] 540), Anti-PD1 (**ab251613**; magenta; Opal[™] 570), Anti-pan Cytokeratin (**ab264485**; red; Opal[™] 620), Anti-EpCAM (**ab225894**; red; Opal[™] 620), Anti-CD8 alpha (**ab251596**; green; Opal[™] 650) and Anti-FOXP3 (**ab96048**; orange; Opal[™] 690). EpCAM and pan-cytokeratin share the same dye and color. Dyes are pseudo-colored for better contrast of the markers.

The immunostaining was performed on a Leica Biosystems BOND[®] MAX instrument with an Opal[™] 6-Plex Detection Kit (NEL821001KT, Akoya Biosciences[®]).

The section was incubated in six rounds of staining; sequentially for **ab251611** (1/750 dilution), **ab219803** (1/250 dilution), **ab251613** (1/750 dilution), **ab264485** (0.5 µg/ml), **ab225894** (1/1250 dilution), **ab251596** (1/1500 dilution) and **ab96048** (10 µg/ml); each using a separate fluorescent tyramide signal amplification system. EDTA based antigen retrieval (Leica Biosystems BOND[®] Epitope Retrieval Solution 2, pH 9.0, 20 minutes) was used in between rounds of tyramide signal amplification to remove the antibody from the previous round, to avoid any cross-reactivity. DAPI (dark blue) was used as a nuclear counter stain.

Microscopy and pseudocoloring of individual Opal[™] dyes was performed using a Vectra 3 Imaging System (Akoya Biosciences[®]).

This data is courtesy of ImmunoAtlas and it can be found [here](#).



Western blot - Anti-FOXP3 antibody [236A/E7] (ab20034)

All lanes : Anti-FOXP3 antibody [236A/E7] (ab20034) at 5 µg/ml

Lane 1 : HEK293T cell lysate

Lane 2 : HEK293T cell lysate overexpressing Human FOXP3

Lane 3 : Human tonsil tissue lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 47 kDa

ab20034 detects Human FOXP3 protein at ~50 kDa in HEK293T cells overexpressing the protein. It also detects FOXP3 in Human tonsil tissue lysate, however this band is significantly weaker in endogenous conditions. Upon higher exposure, weak bands can also be observed in HEK293T cell lysate.

This blot was produced using a 4-12% Bis-Tris gel under the MOPS buffer system. The gel was run at 200V for 60 minutes before being transferred onto a nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour before being incubated with Anti-FOXP3 antibody [236A/E7] (ab20034; 5 microgram per mL) overnight at 4°C. Antibody binding was detected using infrared labelled goat anti-mouse (green; 1:10000) for 1 hour at room temperature before imaging.



Multiplex immunohistochemistry - Anti-FOXP3 antibody [236A/E7] (ab20034)

This image is courtesy of ImmunoAtlas.

Fluorescence multiplex immunohistochemical analysis of Human breast cancer tissue (formalin-fixed paraffin-embedded section).


Merged staining of Anti-PD-L1 ([ab251611](#); cyan; Opal™ 520), Anti-Granzyme B ([ab219803](#); yellow; Opal™ 540), Anti-PD1 ([ab251613](#); magenta; Opal™ 570), Anti-pan Cytokeratin ([ab264485](#); red; Opal™ 620), Anti-EpCAM ([ab225894](#); red; Opal™ 620), Anti-CD8 alpha ([ab251596](#); green; Opal™ 650) and Anti-FOXP3 ([ab96048](#); orange; Opal™ 690). EpCAM and pan-cytokeratin share the same dye and color. Dyes are pseudo-colored for better contrast of the markers.

The immunostaining was performed on a Leica Biosystems BOND® MAX instrument with an Opal™ 6-Plex Detection Kit (NEL821001KT, Akoya Biosciences®).

The section was incubated in six rounds of staining; sequentially for [ab251611](#) (1/750 dilution), [ab219803](#) (1/250 dilution), [ab251613](#) (1/750 dilution), [ab264485](#) (0.5 µg/ml), [ab225894](#) (1/1250 dilution), [ab251596](#) (1/1500 dilution) and [ab96048](#) (10 µg/ml); each using a separate fluorescent tyramide signal amplification system. EDTA based antigen retrieval (Leica Biosystems BOND® Epitope Retrieval Solution 2, pH 9.0, 20 minutes) was used in between rounds of tyramide signal amplification to remove the antibody from the previous round, to avoid any cross-reactivity. DAPI (dark blue) was used as a nuclear counter stain.

Microscopy and pseudocoloring of individual Opal™ dyes was performed using a Vectra 3 Imaging System (Akoya Biosciences®).

This data is courtesy of ImmunoAtlas and it can be found [here](#).



Multiplex immunohistochemistry - Anti-FOXP3 antibody [236A/E7] (ab20034)

This image is courtesy of ImmunoAtlas.

Fluorescence multiplex immunohistochemical analysis of Human breast cancer tissue (formalin-fixed paraffin-embedded section).

Merged staining of Anti-PD-L1 ([ab251611](#); cyan; Opal™ 520), Anti-Granzyme B ([ab219803](#); yellow; Opal™ 540), Anti-PD1 ([ab251613](#); magenta; Opal™ 570), Anti-pan Cytokeratin ([ab264485](#); red; Opal™ 620), Anti-EpCAM ([ab225894](#); red; Opal™ 620), Anti-CD8 alpha ([ab251596](#); green; Opal™ 650) and Anti-FOXP3 ([ab96048](#); orange; Opal™ 690). EpCAM and pan-cytokeratin share the same dye and color. Dyes are pseudo-colored for better contrast of the markers.


The immunostaining was performed on a Leica Biosystems BOND® MAX instrument with an Opal™ 6-Plex Detection Kit (NEL821001KT, Akoya Biosciences®).

The section was incubated in six rounds of staining; sequentially for [ab251611](#) (1/750 dilution), [ab219803](#) (1/250 dilution), [ab251613](#)

(1/750 dilution), **ab264485** (0.5 µg/ml), **ab225894** (1/1250 dilution), **ab251596** (1/1500 dilution) and **ab96048** (10 µg/ml); each using a separate fluorescent tyramide signal amplification system. EDTA based antigen retrieval (Leica Biosystems BOND[®] Epitope Retrieval Solution 2, pH 9.0, 20 minutes) was used in between rounds of tyramide signal amplification to remove the antibody from the previous round, to avoid any cross-reactivity. DAPI (dark blue) was used as a nuclear counter stain.

Microscopy and pseudocoloring of individual Opal[™] dyes was performed using a Vectra 3 Imaging System (Akoya Biosciences[®]).

This data is courtesy of ImmunoAtlas and it can be found [here](#).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXP3 antibody [236A/E7] (ab20034)

Immunohistochemical analysis of paraffin-embedded human tonsil tissue labeling FOXP3 with **ab23004** at 1/500 dilution, followed by a ready to use Goat Anti-Mouse IgG. Counterstained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Goat Anti-Mouse IgG.

Perform heat mediated antigen retrieval using Tris/EDTA buffer (pH 9.0).



Western blot - Anti-FOXP3 antibody [236A/E7] (ab20034)

All lanes : Anti-FOXP3 antibody [236A/E7] (ab20034) at 1/1000 dilution

Lane 1 : HEK-293 (human embryonic kidney) transfected with an empty vector (vector control), containing a GFP-Myc-tag, whole cell lysate


Lane 2 : HEK-293 transfected with FOXP3 (WT) expression vector containing a GFP-Myc-tag, whole cell lysate

Lane 2 : HEK-293 transfected with FOXP3 (WT) expression vector containing a GFP-Myc-tag, whole cell lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 47 kDa

Exposure time: 1 second




Immunohistochemical analysis of paraffin-embedded human thymus tissue labeling FOXP3 with **ab23004** at 1/500 dilution, followed by a ready to use Goat Anti-Mouse IgG. Counterstained with hematoxylin.


Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Goat Anti-Mouse IgG.

Perform heat mediated antigen retrieval using Tris/EDTA buffer (pH 9.0).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXP3 antibody [236A/E7] (ab20034)



(a)




(b)

Immunohistochemical analysis of human large and locally advanced breast cancers staining FOXP3 using ab20034. (a) Low level of FOXP3⁺, CTLA-4⁺ Treg infiltration (b) High level of FOXP3⁺ and CTLA-4⁺ Treg infiltration. (Itu: intratumoral Str: stromal)

This image was generated from the hybridoma version of the product.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXP3 antibody [236A/E7] (ab20034)

This image is from PubMedId: 27777963. Kaewkangsadan Vet al. (2016) Reproduced under the Creative Commons license <https://creativecommons.org/licenses/by/4.0/>




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXP3 antibody

[236A/E7] (ab20034)

Image is courtesy of an AbReview submitted by Jing Ma

ab20034 staining FOXP3 in Cynomolgus Monkey Spleen tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 10% donkey serum for 20 minutes at room temperature; antigen retrieval was by heat mediation using EDTA, pH9.0. Samples were incubated with primary antibody (1/100) for 30 minutes. A Biotin-conjugated Donkey anti-mouse polyclonal (1/2000) was used as the secondary antibody.

This image was generated from the hybridoma version of the product.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXP3 antibody


[236A/E7] (ab20034)

Hermans C. et al PLoS One. 2014 Nov 3;9(11):e111757. doi: 10.1371/journal.pone.0111757. eCollection 2014. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

FoxP3+ cells mainly accumulate centrally.

The formalin-fixed, paraffin-embedded blocks were cut into approximately <math><2\ \mu\text{m}</math> thick slices and mounted on SuperFrost Plus microscope slides (Menzel Gläser, Braunschweig, Germany). After deparaffinization and rehydration, sections were immersed into Dako Target Retrieval solution (Dako North America Inc., Carpinteria, USA), pH 6, 1/10, incubated at 97°C–99°C at 750 Watt for 2x 15 minutes, and allowed to cool to room temperature for 20 minutes. Endogenous peroxidase activity was blocked by 10-minute incubation in 7.5% hydrogen-peroxide solution (Hydroxen Peroxide Solution, Sigma Aldrich Co., Munich, Germany). Immunohistochemical staining for FoxP3 (1/180 dilution; for 60 min) was performed according to standard procedure using MACH-3 mouse alkaline phosphatase polymer detection kit from Biocare Medical Systems (Concord, USA). The slides were incubated with monoclonal mouse antibody. Chromogen Red (Dako North America Inc., Carpinteria, USA) was used as chromogen for FoxP3 staining, and lastly hematoxylin counterstaining was done (Vector Laboratories, Burlingame, USA).

This image was generated from the hybridoma version of the product.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXP3 antibody [236A/E7] (ab20034)

This image is a courtesy of Nicole Schechter

ab20034 staining FOXP3 in human colon tissue section by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). Tissue underwent formaldehyde fixation before heat mediated antigen retrieval and then blocking with 10% serum for 2 hours at 21°C. The primary antibody was diluted 1/50 and incubated with sample for 2 hours at 21°C. An Alexa Fluor®488-conjugated rabbit polyclonal to mouse IgG was used undiluted as secondary antibody.

This image was generated from the hybridoma version of the product.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors