

Anti-Nicastrin antibody ab3444

[4 References](#) [3 图像](#)

概述

产品名称	Anti-Nicastrin抗体
描述	兔多克隆抗体to Nicastrin
宿主	Rabbit
经测试应用	适用于: WB
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide corresponding to Human Nicastrin aa 650-750. Run BLAST with ExPASy Run BLAST with NCBI
阳性对照	WB: T98G, RSC96, mouse neuronal cells, HeLa, IMR-32, NIH/3T3, L6, mouse brain, rat brain
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 99% PBS
纯度	Immunogen affinity purified
Primary antibody说明	Amyloid beta peptide is the major constituent of amyloid plaques in the brains of individuals afflicted with Alzheimer's disease. This peptide is generated from the beta-amyloid precursor protein (beta APP) in a two-step process. The first step involves cleavage of the extracellular, amino-terminal domain of beta APP. Protein cleavage is performed by an aspartyl protease termed beta-secretase (BACE). This enzyme is synthesized as a propeptide that must be modified to the mature and active form by the prohormone convertase, furin. Beta APP cleavage by the mature form of BACE results in the cellular secretion of a segment of beta APP and a

membrane-bound remnant. This remnant is then processed by another protease termed gamma-secretase. Gamma-secretase cleaves an intra-membrane site in the carboxyl-terminal domain of beta APP, thus generating the amyloid beta peptide. Gamma-secretase is believed to be a multi-subunit complex containing presenilin-1 and 2 as central components. Found associated with the presenilins is the transmembrane glycoprotein nicastrin. Nicastrin has been found to bind to the carboxyl-terminus of beta APP and helps to modulate the production of the amyloid beta peptide. Also found in the neurofibrillary lesions associated with Alzheimer's disease is a protein termed Tau. Tau is a neuronal microtubule-associated protein found predominantly on axons. Tau functions to promote tubulin polymerization and stabilize microtubules. Tau, in its hyperphosphorylated form, is the major constituent of paired helical filaments (PHF), which are the building block of neurofibrillary lesions found in brain tissue of Alzheimer's diseased patients.

克隆 多克隆
同种型 IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab3444于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use a concentration of 1 - 2 µg/ml. Detects a band of approximately 110 kDa (predicted molecular weight: 78 kDa).

靶标


功能 Essential subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (beta-amyloid precursor protein). It probably represents a stabilizing cofactor required for the assembly of the gamma-secretase complex.

组织特异性 Widely expressed.

序列相似性 Belongs to the nicastrin family.

细胞定位 Membrane. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

图片



Western blot - Anti-Nicastrin antibody (ab3444)

All lanes : Anti-Nicastrin antibody (ab3444) at 1 µg/ml

Lane 1 : HeLa whole cell extract

Lane 2 : IMR-32 whole cell extract (human neuroblastoma cell line)

Lane 3 : NIH/3T3 whole cell extract (mouse embryo fibroblast cell line)

Lane 4 : L6 whole cell extract (Rat skeletal muscle cell line)

Lane 5 : mouse brain

Lane 6 : rat brain


Lysates/proteins at 60 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG (H+L) Superclonal™ Recombinant Secondary Antibody, HRP at 1/4000 dilution

Developed using the ECL technique.

Predicted band size: 78 kDa



Western blot - Anti-Nicastrin antibody (ab3444)

All lanes : Anti-Nicastrin antibody (ab3444) at 2 µg/ml

Lane 1 : T98G membrane extract (Human glioblastoma cell line)

Lane 2 : RSC96 membrane extract (spontaneously immortalized rat Schwann cell line)

Lysates/proteins at 30 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG (H+L) Superclonal™ Secondary Antibody, HRP conjugate at 1/4000 dilution

Developed using the ECL technique.

Predicted band size: 78 kDa



Western blot detection of Nicastrin on mouse primary cultured neuronal cells using ab3444.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors