


Andrographolide, anti-inflammatory agent ab120636

2 图像


概述

| | |
|-------|--|
| 产品名称 | Andrographolide, anti-inflammatory agent |
| 描述 | Potent anti-inflammatory agent |
| 生物学描述 | Bioactive component of <i>Andrographis paniculata</i> . Potent anti-inflammatory, immunosuppressant, antithrombotic and antiviral agent. Additionally displays antioxidative stress, neuroprotective and antinociceptive activities. |
| 纯度 | > 99% |
| CAS编号 | 5508-58-7 |
| 化学结构 |  |


性能

| | |
|-------|---|
| 化学名称 | (3E,4S)-3-[2-[(1R,4aS,5R,6R,8aS)-Decahydro-6-hydroxy-5-(hydroxymethyl)-5,8a-dimethyl-2-methylene-1-naphthalenyl]ethylidene]dihydro-4-hydroxy-2(3H)-furanone |
| 分子量 | 350.45 |
| 分子式 | C ₂₀ H ₃₀ O ₅ |
| 存放说明 | Store at +4°C. The product can be stored for up to 12 months. |
| 溶解度概述 | Soluble in DMSO to 100 mM and in ethanol to 10 mM |
| 处理 | <p>Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.</p> <p>Refer to SDS for further information</p> <p>Need more advice on solubility, usage and handling? Please visit our frequently asked questions (FAQ) page for more details.</p> |

图片



2D chemical structure image of ab120636, Andrographolide, anti-inflammatory agent



MALME-3M cells were incubated at 37°C for 24h with vehicle control (0 μ M) and 1 μ M andrographolide (ab120636). Increased expression of p53 in MALME-3M cells correlates with an increase in andrographolide concentration, as described in literature.

Whole cell lysates were prepared with RIPA buffer (containing protease inhibitors and sodium orthovanadate), 10 μ g of each were loaded on the gel and the WB was run under reducing conditions. After transfer the membrane was blocked for an hour using 5% BSA before being incubated with [ab26](#) at 5 μ g/ml and [ab8227](#) at 1 μ g/ml overnight at 4°C. Antibody binding was detected using an anti-mouse antibody conjugated to HRP ([ab97040](#)) at 1/10000 dilution and visualised using ECL development solution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team