


Bafilomycin A1, (V)-ATPase inhibitor ab120497

[61 References](#) [4 图像](#)

概述

产品名称	Bafilomycin A1, (V)-ATPase抑制剂
描述	Highly potent, selective (V)-ATPase抑制剂
生物学描述	Highly potent, selective vacuolar (V)-ATPase inhibitor (IC ₅₀ = 500 pM in chromaffin granule membranes). Induces vacuolar deacidification and promotes apoptosis.
CAS编号	88899-55-2
化学结构	

性能

化学名称	(3Z,5E,7R,8S,9S,11E,13E,15S,16R)-8-Hydroxy-16-[(1S,2R,3S)-2-hydroxy-1-methyl-3-[(2R,4R,5S,6R)-tetrahydro-2,4-dihydroxy-5-methyl-6-(1-methylethyl)-2H-pyran-2-yl]butyl]-3,15-dimethoxy-5,7,9,11-tetramethyloxacyclohexadeca-3,5,11,13-tetraen-2-one
分子量	622.84
分子式	C ₃₅ H ₅₈ O ₉
PubChem识别号	6436223
存放说明	Store at -20°C. Store under desiccating conditions. The product can be stored for up to 12 months.
溶解度概述	Soluble in ethanol to 5mM
处理	<p>This product is supplied in one (or more) pack size which is freeze dried. Therefore the contents may not be readily visible, as they can coat the bottom or walls of the vial. Please see our FAQs and information page for more details on handling.</p> <p>Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.</p> <p>Need more advice on solubility, usage and handling? Please visit our frequently asked</p>

[questions \(FAQ\) page](#) for more details.


SMILES

```
O[C@@]1(C[C@@H](O)[C@H](C)[C@H](O1)C(C)C)[C@@H](C)[C@H](O)[C@H](C)C2OC(=O)C(OC)=CC=C(C)[C@@H](C)[C@@H](O)[C@@H](C)CC(C)=CC=C[C@@H]2OC
```

来源


Streptomyces griseus

图片




ab240635 staining SQSTM1/p62 (autophagosome) in control HeLa cells (left panel) and SQSTM1/p62 in HeLa cells treated with 100nM bafilomycin A1 (ab120497) for 18hrs (right panel). The cells were fixed with methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with **ab240635** at 2ug/ml and **ab7291** (Tubulin) at 1/1000 dilution overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (**ab150120**) at 2 µg/ml (shown in red). Nuclear DNA was labelled in blue with DAPI.

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.




2D chemical structure image of ab120497, Bafilomycin A1, (V)-ATPase inhibitor



Immunocytochemistry/ Immunofluorescence -
Bafilomycin A1, (V)-ATPase inhibitor (ab120497)

ab155686 staining SQSTM1/p62 (autophagosome) in control HeLa cells (left panel) and SQSTM1/p62 in HeLa cells treated with 1uM bafilomycin A1 (ab120497) for 18hrs (right panel). The cells were fixed with methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with **ab155686** at 1/500 dilution and **ab7291** (Tubulin) at 1/1000 dilution overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (**ab150120**) at 2 µg/ml (shown in pseudo color red). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunocytochemistry/ Immunofluorescence -
Bafilomycin A1, (V)-ATPase inhibitor (ab120497)

ab109012 staining SQSTM1/p62 (autophagosome) in control HeLa cells (left panel) and SQSTM1/p62 in HeLa cells treated with 1uM bafilomycin A1 (ab120497) for 18hrs (right panel). The cells were fixed with methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with **ab109012** at 5ug/ml and **ab7291** (Tubulin) at 1/1000 dilution overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (**ab150120**) at 2 µg/ml (shown in pseudo color red). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team