

Ro 31-8220 mesylate (Bisindolylmaleimide IX), Protein kinase inhibitor ab120374

[1 References](#) [2 图像](#)

概述

产品名称	Ro 31-8220 mesylate (Bisindolylmaleimide IX), 蛋白kinase抑制剂
描述	蛋白kinase抑制剂
生物学描述	Widely used as a cell-permeable, reversible, competitive protein kinase C inhibitor. Also potently inhibits wide range of kinases including RSK1, RSK2, PRK2, PKC α , PKD1, MSK1, GSK3 β , CDK2-cyclin A and PIM3, as well as PKC α .

Also available in simple stock solutions ([ab146723](#)) - add 1 ml of water to get an exact, ready-to-use concentration.

纯度 > 98%

CAS编号 138489-18-6

化学结构



性能

化学名称	3-[1-[3-(Amidinothio)propyl]-3-indolyl]-4-(1-methyl-3-indolyl)-1 <i>H</i> -pyrrole-2,5-dione methanesulfonate
分子量	553.65
PubChem识别号	11628205
存放说明	Store at -20°C. Store under desiccating conditions. The product can be stored for up to 12 months.
溶解度概述	Soluble in DMSO and organic solvents. Solubility in these solvents is approximately 20 mg/mL.
处理	Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room

temperature for at least 1 hour.

Need more advice on solubility, usage and handling? Please visit our [frequently asked questions \(FAQ\) page](#) for more details.


SMILES

CS(=O)(=O)O.Cn2cc(c1cccc12)C=5C(=O)NC(=O)C=5c4cn(CCCSC(=N)N)c3cccc34

来源


Synthetic

图片



Chemical Structure - Ro 31-8220 mesylate
(Bisindolylmaleimide IX), Protein kinase inhibitor
(ab120374)

2D chemical structure image of ab120374, Ro 31-8220 mesylate
(Bisindolylmaleimide IX), Protein kinase inhibitor



Functional Studies - Ro 31-8220 mesylate
(Bisindolylmaleimide IX), Protein kinase inhibitor
(ab120374)

1.5 x 10⁷ THP-1 cells were incubated with 100 nM GF 109203X ([ab144264](#)) or Ro 31-8220 mesylate (ab120374) for 30 minutes prior to activation with 10 µg x mL⁻¹ PMA (Sigma) for 4 hours. Control cells were left without inhibitors or PMA. Cells were lysed in 1 mL of lysis buffer, and 30 µL were tested for PKC activity (duplicates; +/- SD).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team