

# Human DYNLL1 (PIN) knockout HeLa cell line ab265265

## 2 图像

### 概述

|                             |  |
|-----------------------------|--|
| <b>产品名称</b>                 | 人DYNLL1 (PIN) knockout HeLa cell line  |
| <b>Parental Cell Line</b>   | HeLa   |
| <b>Organism</b>             | Human  |
| <b>Mutation description</b> | Knockout achieved by using CRISPR/Cas9, Homozygous: 1 bp insertion in exon 2   |
| <b>Passage number</b>       | <20  |
| <b>Knockout validation</b>  | Sanger Sequencing, Western Blot (WB)   |
| <b>经测试应用</b>                | <b>适用于:</b> WB   |
| <b>Biosafety level</b>      | 2  |
| <b>常规说明</b>                 | <p><b>Recommended control:</b> Human wild-type HeLa cell line (<a href="#">ab255928</a>). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.</p> <p><b>Cryopreservation cell medium:</b> Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.</p> <p><b>Culture medium:</b> DMEM (High Glucose) + 10% FBS</p> <p><b>Initial handling guidelines:</b> Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.</p> <ol style="list-style-type: none"> <li>1. Thaw the vial in 37°C water bath for approximately 1-2 minutes.</li> <li>2. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution.</li> <li>3. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of <math>2 \times 10^4</math> cells/cm<sup>2</sup>. Seeding density is given as a guide only and should be scaled to align with individual lab schedules.</li> <li>4. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>. Cultures should be monitored daily.</li> </ol> <p><b>Subculture guidelines:</b></p> <p>All seeding densities should be based on cell counts gained by established methods. A guide seeding density of <math>2 \times 10^4</math> cells/cm<sup>2</sup> is recommended.</p> <p>A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.</p> |

Cells should be passaged when they have achieved 80-90% confluence.

This product is subject to limited use licenses from The Broad Institute, ERS Genomics Limited and Sigma-Aldrich Co. LLC, and is developed with patented technology. For full details of the licenses and patents please refer to our [limited use license](#) and [patent pages](#).

We will provide viable cells that proliferate on revival.

## 性能

---

|                              |   |
|------------------------------|---|
| <b>Number of cells</b>       | 1 x 10 <sup>6</sup> cells/vial, 1 mL  |
| <b>Adherent /Suspension</b>  | Adherent  |
| <b>Tissue</b>                | Cervix  |
| <b>Cell type</b>             | epithelial  |
| <b>Disease</b>               | Adenocarcinoma  |
| <b>Gender</b>                | Female  |
| <b>STR Analysis</b>          | Amelogenin X D5S818: 11, 12 D13S317: 12, 13.3 D7S820: 8, 12 D16S539: 9, 10 WWA: 16, 18 TH01: 7 TPOX: 8,12 CSF1PO: 9, 10 |
| <b>Antibiotic resistance</b> | Puromycin 1.00µg/ml   |
| <b>Mycoplasma free</b>       | Yes   |
| <b>存放说明</b>                  | Shipped on Dry Ice. Store in liquid nitrogen.   |
| <b>存储溶液</b>                  | Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether  |

## 靶标

---

|              |   |
|--------------|---|
| <b>功能</b>    | <p>Acts as one of several non-catalytic accessory components of the cytoplasmic dynein 1 complex that are thought to be involved in linking dynein to cargos and to adapter proteins that regulate dynein function. Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. May play a role in changing or maintaining the spatial distribution of cytoskeletal structures.</p> <p>Binds and inhibits the catalytic activity of neuronal nitric oxide synthase.</p> <p>Promotes transactivation functions of ESR1 and plays a role in the nuclear localization of ESR1.</p> <p>Regulates apoptotic activities of BCL2L11 by sequestering it to microtubules. Upon apoptotic stimuli the BCL2L11-DYNLL1 complex dissociates from cytoplasmic dynein and translocates to mitochondria and sequesters BCL2 thus neutralizing its antiapoptotic activity.</p> |
| <b>组织特异性</b> | Ubiquitous.   |
| <b>序列相似性</b> | Belongs to the dynein light chain family.   |
| <b>翻译后修饰</b> | Phosphorylation at Ser-88 appears to control the dimer-monomer transition. According to PubMed:15193260, it is phosphorylated at Ser-88 by PAK1, however, according to PubMed:18650427, the DYNLL1 dimer is not accessible for PAK1 and the phosphorylation could not be demonstrated in vitro.   |
| <b>细胞定位</b>  | Cytoplasm, cytoskeleton. Nucleus. Mitochondrion. Upon induction of apoptosis translocates together with BCL2L11 to mitochondria.  |


## 应用

---

“应用说明”部分下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论 | 说明   |
|----|------|--|
| WB |      | Use at an assay dependent concentration. Predicted molecular weight: 10 kDa. |

## 图片



**All lanes** : Anti-DYNLL1/PIN antibody [EP1660Y] ([ab51603](#)) at 1/1000 dilution

**Lane 1** : Wild-type HeLa cell lysate

**Lane 2** : DYNLL1 knockout HeLa cell lysate

**Lane 3** : Mouse testis tissue lysate

Lysates/proteins at 20 µg per lane.

### Secondary


**All lanes** : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution

**Predicted band size:** 10 kDa

**Observed band size:** 10 kDa

**Lanes 1-3:** Merged signal (red and green). Green - [ab51603](#) observed at 10 kDa. Red - loading control [ab8245](#) observed at 36 kDa.

[ab51603](#) Anti-DYNLL1/PIN antibody [EP1660Y] was shown to specifically react with DYNLL1/PIN in wild-type HeLa cells. Loss of signal was observed when knockout cell line ab265265 (knockout cell lysate [ab257414](#)) was used. Wild-type and DYNLL1/PIN knockout samples were subjected to SDS-PAGE. [ab51603](#) and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Homozygous: 1 bp insertion in exon 2.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors