

Anti-14-3-3 zeta antibody ab51129

★★★★★ [3 Abreviews](#) [30 References](#) [5 图像](#)

概述

产品名称	Anti-14-3-3 zeta抗体
描述	兔多克隆抗体to 14-3-3 zeta
宿主	Rabbit
特异性	<p>ab51129 detects endogenous levels of total 14-3-3 zeta protein.</p> <p>Cross reactivity of ab51129 antibody with other member of the 14-3-3 family:</p> <ul style="list-style-type: none"> -92% identities with 14-3-3 beta/alpha, gamma and eta -77% identities with 14-3-3 teta -69% identities with 14-3-3 epsilon -62% identities with 14-3-3 sigma
经测试应用	适用于: WB, ICC/IF, IHC-P
种属反应性	与反应: Human
免疫原	<p>Synthetic peptide corresponding to Human 14-3-3 zeta. The immunogen is a 13 amino acid synthetic peptide that resides within amino acids 24-73 of the human 14-3-3 zeta.</p> <p>Database link: P63104</p>
阳性对照	WB: 293 cells IHC-P: Human glioblastoma, human breast carcinoma tissue. ICC/IF: HeLa cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: 50% Glycerol, 0.87% Sodium chloride, PBS</p>

	Without Mg+2 and Ca+2
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab51129于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (3)	1/500 - 1/1000. Detects a band of approximately 28 kDa (predicted molecular weight: 28 kDa).
ICC/IF		Use a concentration of 1 µg/ml.
IHC-P		1/50 - 1/100.

靶标

功能	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner.
序列相似性	Belongs to the 14-3-3 family.
翻译后修饰	The delta, brain-specific form differs from the zeta form in being phosphorylated (By similarity). Phosphorylation on Ser-184 by MAPK8; promotes dissociation of BAX and translocation of BAX to mitochondria. Phosphorylation on Ser-58 by PKA; disrupts homodimerization and heterodimerization with YHAE and TP53. This phosphorylation appears to be activated by sphingosine. Phosphorylation on Thr-232; inhibits binding of RAF1.
细胞定位	Cytoplasm. Melanosome. Located to stage I to stage IV melanosomes.

图片




All lanes : Anti-14-3-3 zeta antibody (ab51129)

Lane 1 : Extracts from 293 cells treated with Forskolin (40nM, 30min) with no peptide

Lane 2 : Extracts from 293 cells treated with Forskolin (40nM, 30min) with immunising peptide

Predicted band size: 28 kDa


Observed band size: 28 kDa




ab51129 staining 14-3-3 zeta in Human glioblastoma samples (A) and low grade tumors (B) by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Antigen retrieval was by heat mediation in a citrate buffer. Samples were incubated with primary antibody (1/50) overnight. Nuclei stained blue with hematoxylin. Scale bars = 20 μ m

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-14-3-3 zeta antibody (ab51129)

Simeone P et al., PLoS One 9:e103030 (2014), Fig 7., doi: 10.1371/journal.pone.0103030 Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>




Western blot - Anti-14-3-3 zeta antibody (ab51129)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-14-3-3 zeta antibody (ab51129)

This image shows paraffin-embedded human breast carcinoma tissue stained with ab51129 at a dilution of 1/100. Right hand image: tissue treated with immunising peptide; left hand image: untreated tissue



Immunocytochemistry/ Immunofluorescence - Anti-14-3-3 zeta antibody (ab51129)

ICC/IF image of ab51129 stained HeLa cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab51129, 1µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors