

### Anti-53BP1 antibody [EPR2172(2)] ab175933

敲除验证
重组
RabMAb

★★★★★
[2 Abreviews](#)
[39 References](#)
[15 图像](#)

#### 概述

产品名称	Anti-53BP1抗体[EPR2172(2)]
描述	兔单克隆抗体[EPR2172(2)] to 53BP1
宿主	Rabbit
经测试应用	<b>适用于:</b> WB, IHC-P, ICC/IF, Flow Cyt (Intra)
种属反应性	<b>与反应:</b> Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HepG2 and HeLa cell lysate and human fetal heart and fetal brain tissue lysates, mouse heart and rat heart tissue lysates. IHC-P: human colon, liver carcinoma and tonsil, mouse and rat liver tissues. ICC/IF: HepG2 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR2172(2)
同种型	IgG

## 应用

### The Abpromise guarantee

**Abpromise™** 承诺保证使用ab175933于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000 - 1/5000. Detects a band of approximately 450 kDa (predicted molecular weight: 214 kDa).
IHC-P		1/60 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <a href="#">IHC antigen retrieval protocols</a> .
ICC/IF	★★★★★ (1)	1/100 - 1/250.
Flow Cyt (Intra)		Use at an assay dependent concentration.

## 靶标

功能	May have a role in checkpoint signaling during mitosis. Enhances TP53-mediated transcriptional activation. Plays a role in the response to DNA damage.
疾病相关	Note=A chromosomal aberration involving TP53BP1 is found in a form of myeloproliferative disorder chronic with eosinophilia. Translocation t(5;15)(q33;q22) with PDGFRB creating a TP53BP1-PDGFRB fusion protein.
序列相似性	Contains 2 BRCT domains.
翻译后修饰	Asymmetrically dimethylated on Arg residues by PRMT1. Methylation is required for DNA binding. Phosphorylated at basal level in the absence of DNA damage. Hyper-phosphorylated in an ATM-dependent manner in response to DNA damage induced by ionizing radiation. Hyper-phosphorylated in an ATR-dependent manner in response to DNA damage induced by UV irradiation.
细胞定位	Nucleus. Chromosome > centromere > kinetochore. Associated with kinetochores. Both nuclear and cytoplasmic in some cells. Recruited to sites of DNA damage, such as double strand breaks. Methylation of histone H4 at 'Lys-20' is required for efficient localization to double strand breaks.

## 图片




Immunocytochemistry/ Immunofluorescence - Anti-53BP1 antibody [EPR2172(2)] (ab175933)

Immunocytochemistry/Immunofluorescence analysis of HepG2 cells labelling 53BP1 with purified ab175933 at 1/200. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/1000) were also used.

Control 1: primary antibody (1/200) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/1000).

Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000).



Western blot - Anti-53BP1 antibody [EPR2172(2)] (ab175933)

**Lane 1:** Wild-type HAP1 cell lysate (20 µg)


**Lane 2:** 53BP1 knockout HAP1 cell lysate (20 µg)

**Lane 3:** HeLa cell lysate (40 µg)

**Lane 4:** HepG2 cell lysate (40 µg)


**Lanes 1 - 4:** Merged signal (red and green). Green - ab175933 observed at 350 kDa. Red - loading control, **ab18058**, observed at 124 kDa.

ab175933 was shown to specifically react with 53BP1 when 53BP1 knockout samples were used. Wild-type and 53BP1 knockout samples were subjected to SDS-PAGE. ab175933 and **ab18058** (loading control to Vinculin) were diluted 1/1000 and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat liver tissue labelling 53BP1 with purified ab175933 at a dilution of 1/60. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-53BP1 antibody  
[EPR2172(2)] (ab175933)




ab175933 staining 53BP1 in the human cell line HepG2 (human hepatocellular carcinoma) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde and the sample was incubated with the primary antibody at a dilution of 1/30. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.

Isotype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)

Flow Cytometry (Intracellular) - Anti-53BP1 antibody  
[EPR2172(2)] (ab175933)



**Lane 1:** Wild-type HAP1 cell lysate (20 µg)

**Lane 2:** 53BP1 knockout HAP1 cell lysate (20 µg)

**Lane 3:** HeLa cell lysate (40 µg)


**Lane 4:** HepG2 cell lysate (40 µg)

**Lanes 1 - 4:** Merged signal (red and green).

Green - Target observed at 350 kDa. Red - loading control,

**ab18058**, observed at 124 kDa.

This western blot image is a comparison between ab175933 and a competitor's top cited rabbit polyclonal antibody.



**All lanes :** Anti-53BP1 antibody [EPR2172(2)] (ab175933) at 1/1000 dilution (purified)

**Lane 1 :** Mouse heart tissue lysate

**Lane 2 :** Mouse brain tissue lysate

Lysates/proteins at 10 µg per lane.


### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

**Predicted band size:** 214 kDa

**Observed band size:** 450 kDa

Blocking and dilution buffer: 5% NFDM /TBST.



Anti-53BP1 antibody [EPR2172(2)] (ab175933) at 20 µg (purified)  
+ Rat heart tissue lysate at 20 µg


### Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


**Predicted band size:** 214 kDa

**Observed band size:** 450 kDa

Blocking and dilution buffer: 5% NFDM /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver tissue labelling 53BP1 with purified ab175933 at a dilution of 1/60. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. [ab97051](#), a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



**All lanes :** Anti-53BP1 antibody [EPR2172(2)] (ab175933) at 1/5000 dilution (purified)

**Lane 1 :** Human fetal heart tissue lysate

**Lane 2 :** HeLa whole cell lysate

Lysates/proteins at 20 µg per lane.


### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 214 kDa


**Observed band size:** 450 kDa

Blocking and dilution buffer: 5% NFDM /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-53BP1 antibody [EPR2172(2)] (ab175933)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human liver carcinoma tissue labelling 53BP1 with purified ab175933 at a dilution of 1/60. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



Western blot - Anti-53BP1 antibody [EPR2172(2)] (ab175933)

**All lanes** : Anti-53BP1 antibody [EPR2172(2)] (ab175933) at 1/1000 dilution (unpurified)

**Lane 1** : HepG2 cell lysate


**Lane 2** : Human fetal brain lysate

**Lane 3** : Human fetal heart lysate

Lysates/proteins at 1/10 dilution per lane.

**Predicted band size:** 214 kDa






Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-53BP1 antibody [EPR2172(2)] (ab175933)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human tonsil tissue labelling 53BP1 with unpurified ab175933 at a dilution of 1/100.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-53BP1 antibody [EPR2172(2)] (ab175933)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human colon tissue labelling 53BP1 with unpurified ab175933 at a dilution of 1/100.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-53BP1 antibody [EPR2172(2)] (ab175933)

Immunocytochemistry/Immunofluorescence analysis of HepG2 cells labelling 53BP1 with unpurified ab175933 at a dilution of 1/100.



### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-53BP1 antibody [EPR2172(2)] (ab175933)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors