

Anti-67kDa Laminin Receptor antibody [EPR8469] ab133645

重组 RabMAb

★★★★★ **1 Abreviews** **22 References** **15 图像**

概述

产品名称	Anti-67kDa Laminin Receptor抗体[EPR8469]
描述	兔单克隆抗体[EPR8469] to 67kDa Laminin Receptor
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), WB, IHC-P, ICC/IF, IP
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	K562, HeLa, HepG2, C6, RAW 264.7, PC-12 and NIH/3T3 cell lysates; Human breast and kidney tissues
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR8469
同种型	IgG

应用


The Abpromise guarantee **Abpromise™**承诺保证使用ab133645于以下的经测试应用
“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/20 - 1/1000. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (1)	1/1000 - 1/10000. Predicted molecular weight: 33 kDa.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF		1/100 - 1/250.
IP		1/10 - 1/100.

靶标

功能	Required for the assembly and/or stability of the 40S ribosomal subunit. Required for the processing of the 20S rRNA-precursor to mature 18S rRNA in a late step of the maturation of 40S ribosomal subunits. Also functions as a cell surface receptor for laminin. Plays a role in cell adhesion to the basement membrane and in the consequent activation of signaling transduction pathways. May play a role in cell fate determination and tissue morphogenesis. Acts as a PPP1R16B-dependent substrate of PPP1CA. Also acts as a receptor for several other ligands, including the pathogenic prion protein, viruses, and bacteria.
序列相似性	Belongs to the ribosomal protein S2P family.
翻译后修饰	Acylated. Acylation may be a prerequisite for conversion of the monomeric 37 kDa laminin receptor precursor (37LRP) to the mature dimeric 67 kDa laminin receptor (67LR), and may provide a mechanism for membrane association. Cleaved by stromelysin-3 (ST3) at the cell surface. Cleavage by stromelysin-3 may be a mechanism to alter cell-extracellular matrix interactions.
细胞定位	Cell membrane. Cytoplasm. Nucleus. 67LR is found at the surface of the plasma membrane, with its C-terminal laminin-binding domain accessible to extracellular ligands. 37LRP is found at the cell surface, in the cytoplasm and in the nucleus (By similarity). Co-localizes with PPP1R16B in the cell membrane.

图片



All lanes : Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645) at 1/10000 dilution

Lane 1 : RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) whole cell lysate

Lane 2 : PC-12 (Rat adrenal gland pheochromocytoma cell line) whole cell lysate

Lane 3 : NIH/3T3 (Mouse embryonic fibroblast cell line) whole cell lysate


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


Predicted band size: 33 kDa

Blocking and diluting buffer: 5% NFDM/TBST




ab133645 staining Prohibitin in mouse kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti-rabbit IgG H&L (HRP) [ab97051](#) was used as the secondary antibody at a dilution of 1/500.

Negative control 1: PBS in place of primary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

ab133645 staining 67kDa Laminin Receptor in HeLa (human cervix adenocarcinoma) cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with 4% Paraformaldehyde and permeabilized with 0.1% Triton X-100. Samples were incubated with primary antibody at a dilution of 1/250. A goat anti rabbit IgG (Alexa Fluor® 488) (**ab150077**) was used as the secondary antibody at a dilution of 1/1000. **ab195889** was used as a counterstain for primary antibody ab133645 at 1/2000. DAPI was used as a nuclear counterstain and PBS as a negative control.




Flow Cytometry (Intracellular) - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

ab133645 staining 67kDa Laminin Receptor in the human cell line MCF7 (Human breast adenocarcinoma cell line) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde and the sample was incubated with the primary antibody at a dilution of 1/20. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.

Isoytype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)




Immunoprecipitation - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

ab133645 immunoprecipitating 67kDa Laminin Receptor. 10µg of cell lysate was incubated with primary antibody at a dilution of 1/20 and VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) at a dilution of 1/1000.

Lane 1: K562 (Human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate (10ug)

Lane 2: K562 (Human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate


Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab133645 in K562(Human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

ab133645 staining 67kDa Laminin Receptor in rat kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti-rabbit IgG H&L (HRP) ([ab97051](#)) was used as the secondary antibody at a dilution of 1/500.


Negative control 1: PBS in place of primary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

ab133645 staining 67kDa Laminin Receptor in mouse kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti-rabbit IgG H&L (HRP) **ab97051** was used as the secondary antibody at a dilution of 1/500.


Negative control 1: PBS in place of primary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

ab133645 staining 67kDa Laminin Receptor in human gastric carcinoma tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti-rabbit IgG H&L (HRP) **ab97051** was used as the secondary antibody at a dilution of 1/500.

Negative control 1: PBS in place of primary antibody.



Western blot - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

All lanes : Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645) at 1/2000 dilution

Lane 1 : K562 (Human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate

Lane 2 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 3 : HepG2 (Human liver hepatocellular carcinoma cell line) whole cell lysate

Lane 4 : C6 (Rat glial tumor cell line) whole cell lysate


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)


Predicted band size: 33 kDa

Diluting and blocking buffer: 5% NFDM/TBST



Flow Cytometry (Intracellular) - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

Overlay histogram showing MCF7 cells stained with ab133645 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab133645, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (**ab150077**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1 µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in MCF7 cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



All lanes : Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645) at 1/1000 dilution

Lane 1 : K562 cell lysate

Lane 2 : HeLa cell lysate

Lane 3 : HepG2 cell lysate

Lane 4 : C6 cell lysate

Lane 5 : RAW 264.7 cell lysate

Lane 6 : PC-12 cell lysate


Lane 7 : NIH/3T3 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 33 kDa



Immunohistochemical analysis of paraffin-embedded Human breast tissue labelling 67kDa Laminin Receptor with ab133645 at 1/100 dilution.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labelling 67kDa Laminin Receptor with ab133645 at 1/100 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645)

This image is courtesy of an anonymous Abreview

All lanes : Anti-67kDa Laminin Receptor antibody [EPR8469] (ab133645) at 1/1000 dilution

All lanes : Mouse intestinal mucosa whole tissue lysate

Lysates/proteins at 30 µg per lane.

Secondary

All lanes : HRP-conjugated goat anti-rabbit IgG polyclonal at 1/2500 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 33 kDa

Observed band size: 55 kDa

Exposure time: 1 minute

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-67kDa Laminin Receptor antibody [EPR8469]
(ab133645)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors