

### Anti-ACE2 antibody [EPR24705-45] ab272500

敲除验证
重组
RabMAb

★★★★★
[2 Abreviews](#)
[2 References](#)
[11 图像](#)

#### 概述

产品名称	Anti-ACE2抗体[EPR24705-45]
描述	兔单克隆抗体[EPR24705-45] to ACE2
宿主	Rabbit
经测试应用	<b>适用于:</b> Flow Cyt, IP, WB, IHC-P, ICC/IF <b>不适用于:</b> IHC-Fr
种属反应性	<b>与反应:</b> Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Human kidney and testis tissue lysates; Calu-3 whole cell lysate. IHC-P: Human kidney and testis tissues. ICC/IF: Calu-3, HEK-293T cells transfected with myc-ACE-2; WT CaCo2. Flow cyt: Calu-3 cells. IP: Human testis tissue lysate.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59.94% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR24705-45

同种型

IgG

应用

The Abpromise guarantee

**Abpromise™**承诺保证使用ab272500于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt	★★★★★ (1)	1/500.
IP		1/30.
WB	★★★★★ (1)	1/1000. Detects a band of approximately 120 kDa (predicted molecular weight: 92 kDa).
IHC-P		1/100. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		1/50.

应用说明

Is unsuitable for IHC-Fr.

靶标

功能

Carboxypeptidase which converts angiotensin I to angiotensin 1-9, a peptide of unknown function, and angiotensin II to angiotensin 1-7, a vasodilator. Also able to hydrolyze apelin-13 and dynorphin-13 with high efficiency. May be an important regulator of heart function. In case of human coronaviruses SARS and HCoV-NL63 infections, serve as functional receptor for the spike glycoprotein of both coronaviruses.

组织特异性

Expressed in endothelial cells from small and large arteries, and in arterial smooth muscle cells. Expressed in lung alveolar epithelial cells, enterocytes of the small intestine, Leydig cells and Sertoli cells (at protein level). Expressed in heart, kidney, testis, and gastrointestinal system.

序列相似性

Belongs to the peptidase M2 family.


翻译后修饰

N-glycosylation on Asn-90 may limit SARS infectivity.

细胞定位


Secreted and Cell membrane.

图片



ab272500 staining ACE2 in wild-type Caco2 cells (top panel) and ACE2 knockout Caco2 cells (**ab273731**) (bottom panel). The cells were fixed with 4% paraformaldehyde (10 min) then permeabilized with 0.1% Tween for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab272500 at 10µg/ml concentration and **ab7291** (Mouse monoclonal to alpha Tubulin) at 1/1000 dilution overnight at 4°C followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (**ab150120**) at 2 µg/ml (shown in red). Nuclear DNA was labelled in blue with DAPI.


Image was taken with a confocal microscope (Leica-Microsystems TCS SP8).



Immunohistochemical analysis of paraffin-embedded Human testis tissue labelling ACE2 with ab272500 at 1/100 (5.0 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond® Polymer Refine Detection). Membranous and cytoplasmic staining in human testis. The section was incubated with ab272500 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond® Polymer Refine Detection) .


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-ACE2 antibody [EPR24705-45] (ab272500)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized 293T cells labelling ACE2 with ab272500 at 1/250 (2.0 ug/ml) dilution, followed by [ab150077](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing positive staining in HEK-293T cells transfected with myc-tagged ACE2 expression vector. Myc-Tag Mouse mAb (Alexa Fluor® 647) was used at 1/100 dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is [ab150077](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Immunocytochemistry/ Immunofluorescence - Anti-ACE2 antibody [EPR24705-45] (ab272500)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Calu-3 cells labelling ACE2 with ab272500 at 1/50 (10.0 ug/ml) dilution, followed by [ab150081](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing membranous staining in subsets of Calu-3 cells. **Negative control:** HeLa (PMID: 32926098).

[ab195889](#) Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) (Red) was used at 1/200 dilution. The Nuclear counterstain was DAPI (Blue).


Secondary antibody only control: Secondary antibody is [ab150081](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.



Flow Cytometry - Anti-ACE2 antibody [EPR24705-45] (ab272500)

Flow cytometric analysis of HeLa (Human cervix adenocarcinoma epithelial cell, Left)/ Calu-3 (human lung adenocarcinoma epithelial cell, Right) cells labelling ACE2 with ab272500 at 1/500 dilution (0.1ug). Goat anti rabbit IgG (Alexa Fluor® 647, [ab150079](#)) at 1/2000 dilution was used as the secondary antibody.

**Negative control:** HeLa (PMID: 32926098). Gated on viable cells.



Western blot - Anti-ACE2 antibody [EPR24705-45] (ab272500)

**All lanes :** Anti-ACE2 antibody [EPR24705-45] (ab272500) at 1/1000 dilution

**Lane 1 :** Calu-3 (Human lung adenocarcinoma epithelial cell) whole cell lysate 20

**Lane 2 :** HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate

#### Secondary

**All lanes :** Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 92 kDa


**Observed band size:** 120 kDa

Blocking and diluting buffer and concentration: 5% NFDm/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID:33008593).

**Negative control:** HeLa (PMID:32926098).

Exposure time: 3 minutes



ACE2 was immunoprecipitated from 0.35 mg Human testis lysate with ab272500 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab272500 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP) (**ab131366**) was used at 1/5000 dilution.


**Lane 1:** Human testis lysate 10 ug

**Lane 2:** ab272500 IP in Human testis lysate

**Lane 3:** Rabbit monoclonal IgG (**ab172730**) instead of ab272500 in human testis whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


Exposure time: 32 seconds



Immunohistochemical analysis of paraffin-embedded Human kidney tissue labelling ACE2 with ab272500 at 1/100 (5.0 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Membranous and cytoplasmic staining in human kidney. The section was incubated with ab272500 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND™ RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) .

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Western blot - Anti-ACE2 antibody [EPR24705-45]  
(ab272500)

**All lanes** : Anti-ACE2 antibody [EPR24705-45] (ab272500) at  
1/1000 dilution

**Lane 1** : Human kidney tissue lysate

**Lane 2** : Human testis tissue lysate

**Lane 3** : Human liver tissue lysate

### Secondary

**All lanes** : VeriBlot for IP secondary antibody(HRP)([ab131366](#)) at  
1/1000 dilution

**Predicted band size:** 92 kDa

**Observed band size:** 120 kDa


Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID:10924499, PMID: 32661139).

**Negative control:** liver (PMID:10924499).

Exposure time: 3 minutes





Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ACE2 antibody [EPR24705-45] (ab272500)

Immunohistochemical analysis of paraffin-embedded Human liver tissue labelling ACE2 with ab272500 at 1/100 (5.0 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). **Negative control:** No staining in human liver. The section was incubated with ab272500 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond® Polymer Refine Detection) .

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

#### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-ACE2 antibody [EPR24705-45] (ab272500)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.



## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors