


Anti-Aconitase 2 antibody [EPR8282(B)] ab129069

敲除验证 重组 RabMAb


★★★★★ 8 Abreviews 16 References 6 图像

概述

产品名称	Anti-Aconitase 2抗体[EPR8282(B)]
描述	兔单克隆抗体[EPR8282(B)] to Aconitase 2
宿主	Rabbit
经测试应用	适用于: WB, IHC-P 不适用于: Flow Cyt
种属反应性	与反应: Human 预测可用于: Mouse, Rat 
免疫原	Synthetic peptide within Human Aconitase 2 aa 50-150. The exact sequence is proprietary.
阳性对照	HeLa, K562 and Jurkat cell lysates; Human fetal heart lysate; Human kidney tissue.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"><li>- High batch-to-batch consistency and reproducibility</li><li>- Improved sensitivity and specificity</li><li>- Long-term security of supply</li><li>- Animal-free production</li></ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
解离常数 (K <sub>D</sub> )	K <sub>D</sub> = 2.90 x 10 <sup>-11</sup> M



存储溶液	pH: 7.20
------	----------

	Preservative: 0.01% Sodium azide
	Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture supernatant
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR8282(B)
同种型	IgG

应用

The Abpromise guarantee
Abpromise™承诺保证使用ab129069于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (7)	1/10000 - 1/50000. Detects a band of approximately 85 kDa (predicted molecular weight: 85 kDa).
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.


应用说明

Is unsuitable for Flow Cyt.

靶标

功能	Catalyzes the isomerization of citrate to isocitrate via cis-aconitate.
通路	Carbohydrate metabolism; tricarboxylic acid cycle; isocitrate from oxaloacetate: step 2/2.
序列相似性	Belongs to the aconitase/IPM isomerase family.
细胞定位	Mitochondrion.

图片



Western blot - Anti-Aconitase 2 antibody  
[EPR8282(B)] (ab129069)

**Lane 1:** Wild-type HAP1 whole cell lysate (20 µg)


**Lane 2:** Aconitase 2 knockout HAP1 whole cell lysate (20 µg)

**Lane 3:** HeLa whole cell lysate (20 µg)

**Lane 4:** Jurkat whole cell lysate (20 µg)

**Lanes 1 - 4:** Merged signal (red and green). Green - ab129069 observed at 90 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

ab129069 was shown to specifically react with Aconitase 2 in wild-type HAP1 cells as signal was lost in Aconitase 2 knockout cells. Wild-type and Aconitase 2 knockout samples were subjected to SDS-PAGE. ab129069 and **ab9484** (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/10000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Aconitase 2 antibody  
[EPR8282(B)] (ab129069)

**All lanes :** Anti-Aconitase 2 antibody [EPR8282(B)] (ab129069) at 1/10000 dilution

**Lane 1 :** HeLa cell lysate

**Lane 2 :** K562 cell lysate

**Lane 3 :** Human fetal heart tissue lysate


**Lane 4 :** Jurkat cell lysate

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes :** Goat anti-rabbit HRP-conjugated at 1/2000 dilution


**Predicted band size:** 85 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aconitase 2 antibody [EPR8282(B)] (ab129069)

ab129069, at 1/100 dilution, staining Aconitase 2 in paraffin-embedded Human kidney tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Western blot - Anti-Aconitase 2 antibody [EPR8282(B)] (ab129069)

This image is courtesy of an abreview submitted by Ruma Raha-Chowdhury, University Of Cambridge, United Kingdom

**All lanes :** Anti-Aconitase 2 antibody [EPR8282(B)] (ab129069) at 1/1000 dilution

**Lane 1 :** 1 Month old Mouse Liver

**Lanes 2-3 :** 3 Month old Mouse Liver

**Lane 4 :** 6 Month old Mouse Liver

#### Secondary

**All lanes :** Dako Pig polyclonal to anti-rabbit (HRP) at 1/3000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 85 kDa

**Exposure time:** 1 minute

**Blocking:** 5% Milk at 24°C for 1 Hour



OI-RD Scanning - Anti-Aconitase 2 antibody  
[EPR8282(B)] (ab129069)

Equilibrium dissociation constant ( $K_D$ )

Learn more about  $K_D$

[Click here to learn more about  \$K\_D\$](#)

Why choose a  
recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Aconitase 2 antibody [EPR8282(B)] (ab129069)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors