# abcam

# Product datasheet

# Anti-ADAM15 antibody [EPR5619] - BSA and Azide free ab232389



RabMAb

3 图像

#### 概述

产品名称 Anti-ADAM15抗体[EPR5619] - BSA and Azide free

描述 兔单克隆抗体[EPR5619] to ADAM15 - BSA and Azide free

宿主 Rabbit

**适用于:** WB, IHC-P

种属反应性 与反应: Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 IHC-P: Human colon tissue.

常规说明 ab232389 is the carrier-free version of ab124698.

Our <u>carrier-free</u> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our <u>conjugation kits</u> for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**<sup>®</sup> **patents**.

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with

1

### 性能

形式 Liquid

**存放说明** Shipped at 4°C. Store at +4°C. Do Not Freeze.

**存储溶液** pH: 7.2

Constituent: PBS

纯**度** Protein A purified

**克隆** 单克隆

**克隆编号** EPR5619

同种型 IgG

#### 应用

## The Abpromise guarantee Abpromise™承诺保证使用ab232389于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use at an assay dependent concentration. Predicted molecular weight: 93 kDa.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

#### 靶标

功能

Active metalloproteinase with gelatinolytic and collagenolytic activity. Plays a role in the wound healing process. Mediates both heterotypic intraepithelial cell/T-cell interactions and homotypic T-cell aggregation. Inhibits beta-1 integrin-mediated cell adhesion and migration of airway smooth muscle cells. Suppresses cell motility on or towards fibronectin possibly by driving alpha-v/beta-1 integrin (ITAGV-ITGB1) cell surface expression via ERK1/2 inactivation. Cleaves E-cadherin in response to growth factor deprivation. Plays a role in glomerular cell migration. Plays a role in pathological neovascularization. May play a role in cartilage remodeling. May be proteolytically processed, during sperm epididymal maturation and the acrosome reaction. May play a role in sperm-egg binding through its disintegrin domain.

组织特异性

Expressed in colon and small intestine. Expressed in airway smooth muscle and glomerular mesangial cells (at protein level). Ubiquitously expressed. Overexpressed in atherosclerotic lesions. Constitutively expressed in cultured endothelium and smooth muscle. Expressed in chondrocytes. Expressed in airway smooth muscle and glomerular mesangial cells.

序列相似性

Contains 1 disintegrin domain.
Contains 1 EGF-like domain.

Contains 1 peptidase M12B domain.

The cytoplasmic domain interacts with endophilin I and sorting nexin 9.

Disintegrin domain binds to integrin alphaV-beta3.

The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-

peptide release activates the enzyme.

The precursor is cleaved by a furin endopeptidase.

 $\label{prop:phosphorylation} Phosphorylation\ increases\ association\ with\ PTKs.$ 

Endomembrane system. Cell junction > adherens junction. Cell projection > cilium > flagellum. Cytoplasmic vesicle > secretory vesicle > acrosome. The majority of the protein is localized in a perinuclear compartment which may correspond to the trans-Golgi network or the late endosome. The pro-protein is the major detectable form on the cell surface, whereas the majority of the


protein in the cell is processed.

### 图片

结构域

翻译后修饰

细胞定位




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ADAM15 antibody

[EPR5619] - BSA and Azide free (ab232389)

**ab124698**, at 1/250 dilution, staining ADAM15 in paraffinembedded Human colon tissue by immunohistochemistry. This image was produced using unpurified antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab124698).




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ADAM15 antibody

[EPR5619] - BSA and Azide free (ab232389)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human colon tissue sections labeling ADAM15 with purified <a href="mailto:ab124698">ab124698</a> at 1:800 dilution. Heat mediated antigen retrieval was performed using <a href="mailto:ab93684">ab93684</a> (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use). PBS instead of the primary antibody was used as the negative control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab124698</u>).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

# Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors