abcam

Product datasheet

Alexa Fluor® 405 Anti-ERK1 + ERK2 antibody [EPR17526] ab210044

重组 RabMAb

2图像

概述			
产品名称	Alexa Fluor® 405荧 光 Anti-ERK1 + ERK2 抗体 [EPR17526]		
描述	Alexa Fluor® 405荧 光兔 单 克隆抗体 [EPR17526] to ERK1 + ERK2		
宿主	Rabbit		
偶 联物	Alexa Fluor® 405. Ex: 402nm, Em: 421nm		
经测试应 用	适用于: Flow Cyt (Intra)		
种属反 应性	与反应: Human		
	预测可用于: Mouse, Rat 🛛 📤		
免疫原	Recombinant fragment within Human ERK1 + ERK2 aa 150 to the C-terminus. The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please contact our Scientific Support team to discuss your requirements. Database link: P28482 Run BLAST with		
阳性 对照	Flow Cyt (intra): A431 cells.		
性能			
形式	Liquid		
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C. Store In the Dark.		
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: 1% BSA, 30% Glycerol (glycerin, glycerine), PBS		
纯 度	Protein A purified		
克隆	单 克隆		
克隆 编号	EPR17526		
同种型	lgG		

应用

The Abpromise guarantee


Abpromise™承诺保证使用ab210044于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/500.

靶 标	
功能	Involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK1. Phosphorylates EIF4EBP1; required for initiation of translation. Phosphorylates microtubule- associated protein 2 (MAP2). Phosphorylates SPZ1 (By similarity). Phosphorylates heat shock factor protein 4 (HSF4) and ARHGEF2. Acts as a transcriptional repressor. Binds to a [GC]AAA[GC] consensus sequence. Repress the expression of interferon gamma-induced genes. Seems to bind to the promoter of CCL5, DMP1, IFIH1, IFITM1, IRF7, IRF9, LAMP3, OAS1, OAS2, OAS3 and STAT1. Transcriptional activity is independent of kinase activity.
序列相似性	Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily. subfamily. Contains 1 protein kinase domain.
结 构域	The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP kinases.
翻 译后 修 饰	Dually phosphorylated on Thr-185 and Tyr-187, which activates the enzyme. Dephosphorylated by PTPRJ at Tyr-187.
细 胞定位	Nucleus.
形式	Mainly expressed in the cytoplasm and only localizes to the nucleus with treatment.

图片



Why choose a recombinant antibody? Research with Long-term and scalable supply confidence Consistent and Recombinant reproducible results technology Success from the Ethical standards first experiment compliant Confirmed Animal-free production specificity Alexa Fluor® 405 Anti-ERK1 + ERK2 antibody [EPR17526] (ab210044)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.cn/abpromise</u> or contact our technical team.

Overlay histogram showing A431 cells stained with ab210044 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Triton X-100 for 15 min, blocked with Image-iT® FX Signal Enhancer for 30 min at 22°C and then blocked with 1x PBS / 10% normal goat serum for 1 hr at 22°C. Cells were then incubated with the antibody (ab210044, 1/500 dilution) for 30 min at 22°C.

Isotype control antibody (black line) was rabbit IgG (monoclonal) Alexa Fluor® 405 (<u>ab208150</u>) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 50mW violet laser (405nm) and 450/50 bandpass filter.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors