abcam

Product datasheet

Alexa Fluor® 488 Anti-BST2/Tetherin antibody [EPR23597-266] ab275192



1 图像

概述

产品名称 Alexa Fluor® 488荧光Anti-BST2/Tetherin抗体[EPR23597-266]

描述 Alexa Fluor® 488荧光兔单克隆抗体[EPR23597-266] to BST2/Tetherin

宿主 Rabbit

偶联物 Alexa Fluor® 488. Ex: 495nm, Em: 519nm

经测试应用适用于: Flow Cyt种属反应性与反应: Mouse

免疫原 Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

阳性对照 Flow cyt: C57 BL/6 mouse splenocytes.

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**[®] **patents**.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor[®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor[®] dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or **outlicensing@thermofisher.com**.

1

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle. Store In the Dark.

存储溶液 pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, 68.98% PBS

纯**度** Protein A purified

克隆 单克隆

克隆编号 EPR23597-266

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab275192于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		1/500.

靶标

功能 May be involved in the sorting of secreted proteins (By similarity). May be involved in pre-B-cell

growth. Antiretroviral defense protein, that blocks release of retrovirus from the cell surface.

Depleted unpon HIV-1 infection by viral VPU protein through 20S proteasome degradation.

Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) through ubiquitination and subsequent degradation. May play a role in B-cell activation in rheumatoid

arthritis.

组织特异性 Predominantly expressed in liver, lung, heart and placenta. Lower levels in pancreas, kidney,

skeletal muscle and brain. Overexpressed in multiple myeloma cells. Highly expressed during B-cell development, from pro-B precursors to plasma cells. Highly expressed on T-cells, monocytes,

NK cells and dendritic cells (at protein level).

序列相似性 Belongs to the tetherin family.

结构域 The extracellular coiled coil domain is important for virus retention at the cell surface and

prevention of virus spreading.


翻译后修饰 Monoubiquitinated by KSHV E3 ubiquitin-protein ligase K5, leading to its targeting to late

endosomes and degradation.

细胞定位 Golgi apparatus > trans-Golgi network. Cell membrane. Cell membrane. Late endosome.

Targeted to late endosomes upon KSHV infection and subsequent ubiquitination. Targeted to the

trans-Golgi network by viral VPU protein.



Flow cytometry staining of C57 BL/6 mouse splenocytes with ab275192 (right) or Rabbit IgG (monoclonal) Alexa Fluor $^{@}$ 488 (ab199091) isotype control (left). Cells were incubated for 30 min on ice in 1x PBS containing 10 μ /ml anti-CD16/CD32 and 10% normal goat serum to block FC receptors and nonspecific protein-protein interaction followed by the antibody (ab275192) or Rabbit IgG (monoclonal) Alexa Fluor 488 $^{@}$ (ab199091) isotype (1 x 10 6 in 100 μ l at 1 μ g/ml (1/500)) for 30 min on ice. The cells were simultaneously stained with CD45R. Acquisition of >30000 events were collected using a 50 mW Blue laser (488nm) and 530/30 bandpass filter. Events were gated on live single cells.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors