abcam

Product datasheet

Alexa Fluor® 488 Anti-GAPDH antibody [EPR16891] ab201768



重组 RabMAb

2 图像

概述

免疫原

产品名称 Alexa Fluor® 488荧光Anti-GAPDH抗体[EPR16891]

描述 Alexa Fluor® 488荧光兔单克隆抗体[EPR16891] to GAPDH

宿主 Rabbit

偶联物 Alexa Fluor® 488. Ex: 495nm, Em: 519nm

经测试应用 适用于: ICC/IF 种属反应性 与反应: Human

预测可用于: Mouse, Rat, Chicken, Fish, Monkey, Zebrafish, Xenopus tropicalis

Recombinant fragment within Mouse GAPDH aa 100 to the C-terminus. The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the

immunogen is needed to determine the suitability of the antibody for your needs, please contact

our Scientific Support team to discuss your requirements.

Database link: P16858

Run BLAST with Run BLAST with

阳性对照 ICC/IF: HeLa cells.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit 常规说明 monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

> Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

存储溶液 pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA

纯**度** Protein A purified

克隆 单克隆

克隆编号 EPR16891

同种型 lgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab201768于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/200.

靶标

功能 Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing

a role in glycolysis and nuclear functions, respectively. Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis. Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D-glyceraldehyde 3-

phosphate (G3P) into 3-phospho-D-glyceroyl phosphate.

通路 Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 1/5.

序列相似性 Belongs to the glyceraldehyde-3-phosphate dehydrogenase family.

翻译后修饰 S-nitrosylation of Cys-152 leads to interaction with SIAH1, followed by translocation to the

nucleus.


ISGylated.

细**胞定位** Cytoplasm > cytosol. Nucleus. Cytoplasm > perinuclear region. Membrane. Translocates to the

nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization

signal (By similarity). Postnuclear and Perinuclear regions.


图片



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-GAPDH antibody [EPR16891] (ab201768)

ab201768 staining GAPDH in HeLa cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab201768 at 1/200 dilution (shown in green) and <u>ab195889</u>, Mouse monoclonal to alpha Tubulin (Alexa Fluor[®] 594), at $2\mu g/ml$ (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors