abcam

Product datasheet

Alexa Fluor® 488 Anti-Nrf2 (phospho S40) antibody [EP1809Y] ab237464

重组 RabMAb

2 图**像**

概述	
产品名称	Alexa Fluor® 488荧 光 Anti-Nrf2 (phospho S40) 抗体 [EP1809Y]
描述	Alexa Fluor® 488荧 光兔 单 克隆抗体 [EP1809Y] to Nrf2 (phospho S40)
宿主	Rabbit
偶 联物	Alexa Fluor® 488. Ex: 495nm, Em: 519nm
经 测 试应 用	适用于: ICC/IF
种属反应性	与反应: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. (Peptide available as ab133404)
阳性 对照	ICC/IF: HepG2 cells
常规说明	 This product is a recombinant monoclonal antibody, which offers several advantages including: High batch-to-batch consistency and reproducibility Improved sensitivity and specificity Long-term security of supply Animal-free production For more information <u>see here</u>. Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <u>RabMAb[®] patents</u>.
	Alexa Fluor [®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor [®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor [®] dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor [®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

outlicensing@thermofisher.com.

Rat: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.

性能	
形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. Stable for 12 months at -20°C. Store In the Dark.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS
纯 度	Protein A purified
克隆	单 克隆
克 隆 编号	EP1809Y
同种型	lgG


应用

The Abpromise guarantee Abpromise ™承诺保证使用ab237464于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/200. This product gave a positive signal in HepG2 cells fixed with 4% formaldehyde (10 min)


靶标	
功能	Transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. Important for the coordinated up-regulation of genes in response to oxidative stress. May be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the beta-globin locus control region.
组织 特异性	Widely expressed. Highest expression in adult muscle, kidney, lung, liver and in fetal muscle.
序列相似性	Belongs to the bZIP family. CNC subfamily. Contains 1 bZIP domain.
结 构域	Acidic activation domain in the N-terminus, and DNA binding domain in the C-terminus.
翻 译 后修 饰	Phosphorylation of Ser-40 by PKC in response to oxidative stress dissociates NFE2L2 from its cytoplasmic inhibitor KEAP1, promoting its translocation into the nucleus.
细 胞定位	Cytoplasm > cytosol. Nucleus. Cytosolic under unstressed conditions, translocates into the nucleus upon induction by electrophilic agents.



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-Nrf2 (phospho S40) antibody [EP1809Y] (ab237464) ab237464 staining Nrf2 (phospho S40) in HepG2 cells +/- Lambda Phosphatase (2 hours at 31°C). The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h.

The cells were then incubated overnight at +4°C with ab237464 at 1/200 dilution (shown in green) and **ab195889**, Mouse monoclonal to alpha Tubulin (Alexa Fluor[®] 594), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.cn/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors