abcam

Product datasheet

Alexa Fluor® 488 Anti-PKC delta antibody [EPR17075] ab206282





RabMAb

1 References 2 图像

概述

产品名称 Alexa Fluor® 488荧光Anti-PKC delta抗体[EPR17075]

Alexa Fluor® 488荧光兔单克隆抗体[EPR17075] to PKC delta 描述

宿主 Rabbit

偶联物 Alexa Fluor® 488. Ex: 495nm, Em: 519nm

适用干: ICC/IF 经测试应用 种属反应性 与反应: Human

预测可用于: Mouse, Rat 🔷

Recombinant fragment within Mouse PKC delta aa 500 to the C-terminus. The exact sequence is 免疫原

proprietary.

Database link: P28867

ICC/IF: HeLa cells 阳性对照

常规说明 This conjugated antibody has been KO validated based on the results obtained with the

unconjugated clone: Anti-PKC delta antibody [EPR17075] (ab182126).

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc., a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or

prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

存储溶液 pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 1% BSA, PBS, 30% Glycerol (glycerin, glycerine)

纯**度** Protein A purified

克隆 单克隆

克隆编号 EPR17075

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab206282于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/50. This product gave a positive signal in HeLa cells fixed with 100% methanol (5 min).

靶标

功能 This is calcium-independent, phospholipid-dependent, serine- and threonine-specific enzyme.

PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters, a class of tumor promoters. May play a role in antigen-dependent control of B-cell function. Phosphorylates MUC1 in the C-terminal and

regulates the interaction between MUC1 and beta-catenin.

序列相似性 Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily.

Contains 1 AGC-kinase C-terminal domain.

Contains 1 C2 domain.

Contains 2 phorbol-ester/DAG-type zinc fingers.

Contains 1 protein kinase domain.

结构域 The C1 domain, containing the phorbol ester/DAG-type region 1 (C1A) and 2 (C1B), is the

diacylglycerol sensor.


The C2 domain is a non-calcium binding domain. It binds proteins containing phosphotyrosine in

a sequence-specific manner.

细胞定位

Cytoplasm. Membrane.


图片



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-PKC delta antibody [EPR17075] (ab206282) ab206282 staining PKC delta in HeLa cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab206282 at 1/50 dilution (shown in green) and ab195889, Mouse monoclonal to alpha Tubulin (Alexa Fluor[®] 594), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Phosphorylated on Thr-507, within the activation loop. Autophosphorylated and/or phosphorylated.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors