abcam

Product datasheet

Alexa Fluor® 647 Anti-BST2/Tetherin antibody [EPR23597-266] ab275193

重组

1 图**像**

概述		
产 品名称	Alexa Fluor® 647荧光Anti-BST2/Tetherin 抗体 [EPR23597-266]	
描述	Alexa Fluor® 647荧 光兔 单 克隆抗体[EPR23597-266] to BST2/Tetherin	
宿主	Rabbit	
偶联物	Alexa Fluor® 647. Ex: 652nm, Em: 668nm	
经测试应 用	适用于: Flow Cyt	
种属反应性	与反应: Mouse	
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.	
阳性 对照	Flow cyt: C57 BL/6 mouse splenocytes.	
常规说明		

性能	
形式	Liquid
存 放 说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. Store In the Dark.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, 68.98% PBS
纯 度	Protein A purified
克隆	单 克隆
克隆 编号	EPR23597-266
同种型	lgG

应用


The Abpromise guarantee

Abpromise™承诺保证使用ab275193于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		1/500.

靶 标	
功能	May be involved in the sorting of secreted proteins (By similarity). May be involved in pre-B-cell growth. Antiretroviral defense protein, that blocks release of retrovirus from the cell surface. Depleted unpon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) through ubiquitination and subsequent degradation. May play a role in B-cell activation in rheumatoid arthritis.
组织 特异性	Predominantly expressed in liver, lung, heart and placenta. Lower levels in pancreas, kidney, skeletal muscle and brain. Overexpressed in multiple myeloma cells. Highly expressed during B-cell development, from pro-B precursors to plasma cells. Highly expressed on T-cells, monocytes, NK cells and dendritic cells (at protein level).
序列相似性	Belongs to the tetherin family.
结 构域	The extracellular coiled coil domain is important for virus retention at the cell surface and prevention of virus spreading.
翻 译 后修 饰	Monoubiquitinated by KSHV E3 ubiquitin-protein ligase K5, leading to its targeting to late endosomes and degradation.
细胞定位	Golgi apparatus > trans-Golgi network. Cell membrane. Cell membrane. Late endosome. Targeted to late endosomes upon KSHV infection and subsequent ubiquitination. Targeted to the trans-Golgi network by viral VPU protein.



Flow cytometry staining of C57 BL/6 mouse splenocytes with ab275193 (right) or Rabbit IgG (monoclonal) Alexa Fluor[®] 647 isotype (left). Cells were incubated for 30 min on ice in 1x PBS containing 10 μ g/mL anti CD16/CD32 and 10 % normal goat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab275193) or Rabbit IgG (monoclonal) Alexa Fluor[®] 647 isotype (1x 10⁶ in 100 μ L at 1 μ g/mL (1/500)) for 30 min on ice. The cells were simultaneously stained with CD45R.

Acquisition of >30000 events were collected using a 40 mW Red laser (638nm) and 660/10 bandpass filter. Events were gated on live single cells.

Flow Cytometry - Alexa Fluor® 647 Anti-BST2/Tetherin antibody [EPR23597-266] (ab275193)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.cn/abpromise</u> or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors