

Anti-ALIX antibody [EPR23653-32] ab275377

敲除验证
重组
RabMAb

★★★★★
[1 Abreviews](#)
[11 References](#)
[10 图像](#)


概述

产品名称	Anti-ALIX抗体[EPR23653-32]
描述	兔单克隆抗体[EPR23653-32] to ALIX
宿主	Rabbit
经测试应用	适用于: WB, ICC/IF, Flow Cyt (Intra), IP 不适用于: IHC-P
种属反应性	与反应: Mouse, Rat, Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: C6, RAW 264.7, PC-12, NIH/3T3, K-562, HEK-293, HeLa, HCT116, MCF7 and Jurkat whole cell lysates; Human brain tissue lysate; Mouse brain tissue lysate; Rat brain tissue lysate. ICC/IF: NIH/3T3 and HeLa cells. Flow Cyt (intra): NIH/3T3 and HeLa cells. IP: K562 whole cell lysate.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.01% Sodium azide Constituents: 59.94% PBS, 40% Glycerol, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR23653-32

同种型	IgG	
应用		
<div>The Abpromise guarantee</div> <div>Abpromise™承诺保证使用ab275377于以下的经测试应用</div> <div>“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。</div>		
应用	Ab评论	说明
WB	★★★★★ (1)	1/1000. Detects a band of approximately 80, 90, 100 kDa (predicted molecular weight: 96 kDa).
ICC/IF		1/5000.
Flow Cyt (Intra)		1/500.
IP		1/30.
应用说明	Is unsuitable for IHC-P.	
靶标		
功能	Class E VPS protein involved in concentration and sorting of cargo proteins of the multivesicular body (MVB) for incorporation into intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome. Binds to the phospholipid lysobisphosphatidic acid (LBPA) which is abundant in MVBs internal membranes. The MVB pathway appears to require the sequential function of ESCRT-O, -I,-II and -III complexes. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and enveloped virus budding (HIV-1 and other lentiviruses). Appears to be an adapter for a subset of ESCRT-III proteins, such as CHMP4, to function at distinct membranes. Required for completion of cytokinesis. Involved in HIV-1 virus budding. Can replace TSG101 it its role of supporting HIV-1 release; this function implies the interaction with CHMP4B. May play a role in the regulation of both apoptosis and cell proliferation.	
序列相似性	Contains 1 BRO1 domain.	
细胞定位	Cytoplasm > cytosol. Melanosome. Cytoplasm > cytoskeleton > centrosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV. Colocalized with CEP55 in the midbody during cytokinesis. Colocalized with CEP55 at centrosomes of non-dividing cells.	
图片		



All lanes : Anti-ALIX antibody [EPR23653-32] (ab275377) at 1/1000 dilution

Lane 1 : Wild-type HEK-293 cell lysate

Lane 2 : PDCD6IP knockout HEK-293 cell lysate

Lane 3 : HeLa cell lysate

Lane 4 : Human Spleen tissue lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 96 kDa

Observed band size: 96 kDa


Lanes 1 -4: Merged signal (red and green). Green - ab275377 observed at 96 kDa. Red - loading control **ab8245** (Mouse anti-GAPDH antibody [6C5]) observed at 37 kDa.

ab275377 was shown to react with PDCD6IP in wild-type HEK-293T cells in Western blot with loss of signal observed in PDCD6IP knockout cell line **ab260864** (PDCD6IP knockout cell lysate **ab261656**). Wild-type HEK-293T and PDCD6IP knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 5 % milk in TBS-T (0.1 % Tween®) before incubation with ab275377 and **ab8245** (Mouse anti-GAPDH antibody [6C5]) overnight at 4 °C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 h at room temperature before imaging.



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HeLa (human epithelial cell line from cervix adenocarcinoma) cells labelling ALIX with ab275377 at 1/5000 dilution, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488, [ab150077](#)) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in HeLa cells is observed. Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594, [ab195889](#)) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488, [ab150077](#)) at 1/1000 dilution.



All lanes : Anti-ALIX antibody [EPR23653-32] (ab275377) at 1/1000 dilution

Lane 1 : Mouse brain tissue lysate

Lane 2 : Rat brain tissue lysate

Lane 3 : C6 (rat glial tumor glial cell) whole cell lysate

Lane 4 : RAW264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate

Lane 5 : PC-12 (rat adrenal gland pheochromocytoma) whole cell lysate

Lane 6 : NIH/3T3 (mouse embryonic fibroblast) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated ([ab97051](#)) at 1/100000 dilution


Predicted band size: 96 kDa

Observed band size: 100,80,90 kDa

Blocking and diluting buffer and concentration: 5% NFDm/TBST.

The expression pattern is consistent with what has been described in the literature (PMID: 24834918, 26935291, 28322231).


Exposure time: Lane1-2: 10 seconds Lane3-6: 8 seconds.



Flow Cytometry (Intracellular) - Anti-ALIX antibody
[EPR23653-32] (ab275377)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol permeabilized HeLa (human cervix adenocarcinoma epithelial cell line) cells labelling ALIX with ab275377 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (**ab172730**) isotype control (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) at 1/2000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-ALIX antibody
[EPR23653-32] (ab275377)

ALIX was immunoprecipitated from 0.35 mg K-562 (human chronic myelogenous leukemia lymphoblast cell line) whole cell lysate with ab275377 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab275377 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used at 1/5000 dilution.


Lane 1: K-562 whole cell lysate 10 ug

Lane 2: ab275377 IP in K-562 whole cell lysate

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab275377 in K-562 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 seconds



Western blot - Anti-ALIX antibody [EPR23653-32] (ab275377)

Anti-ALIX antibody [EPR23653-32] (ab275377) at 1/1000 dilution + Human brain tissue lysate at 20 µg

Secondary

VeriBlot for IP secondary antibody(HRP)([ab131366](#)) at 1/1000 dilution


Predicted band size: 96 kDa

Observed band size: 100,80,90 kDa

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The expression pattern is consistent with what has been described in the literature (PMID: 24834918, 26935291, 28322231).


Exposure time: 10 seconds.



Flow Cytometry (Intracellular) - Anti-ALIX antibody [EPR23653-32] (ab275377)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol permeabilized NIH/3T3 (mouse embryonic fibroblast cell line) cells labelling ALIX with ab275377 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG ([ab172730](#)) isotype control (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).

A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) at 1/2000 dilution was used as the secondary antibody.



Western blot - Anti-ALIX antibody [EPR23653-32]
(ab275377)

All lanes : Anti-ALIX antibody [EPR23653-32] (ab275377) at 1/1000 dilution

Lane 1 : K-562 (human chronic myelogenous leukemia lymphoblast) whole cell lysate

Lane 2 : HEK-293 (human embryonic kidney epithelial cell) whole cell lysate

Lane 3 : HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate

Lane 4 : HCT116 (human colorectal carcinoma epithelial cell) whole cell lysate

Lane 5 : MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysate

Lane 6 : Jurkat (human T cell leukemia T lymphocyte) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated ([ab97051](#)) at 1/100000 dilution


Predicted band size: 96 kDa

Observed band size: 90-100 kDa

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The expression pattern is consistent with what has been described in the literature (PMID: 24834918, 26935291, 28322231).

Exposure time: 10 seconds.



Immunocytochemistry/ Immunofluorescence - Anti-ALIX antibody [EPR23653-32] (ab275377)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (mouse embryo fibroblast cell line) cells labelling ALIX with ab275377 at 1/5000 dilution, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488, [ab150077](#)) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in NIH/3T3 cells. Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594, [ab195889](#)) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is [ab150077](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-ALIX antibody [EPR23653-32] (ab275377)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors