

Anti-alpha Actinin 4 antibody [EPR2533(2)] ab108198

敲除验证
重组
RabMAb

★★★★★
9 Abreviews
26 References
11 图像

概述

产品名称	Anti-alpha Actinin 4抗体[EPR2533(2)]
描述	兔单克隆抗体[EPR2533(2)] to alpha Actinin 4
宿主	Rabbit
经测试应用	适用于: WB, IP, IHC-P, ICC/IF, Flow Cyt (Intra)
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Human skeletal muscle, A431, Mouse brain, Rat brain, Rat Heart, HAP1 and MFC7 Lysates Flow Cyt: HeLa IHC-P: Human breast carcinoma, Mouse and Rat liver tissues ICC/IF: HAP1 cells (HAP1-ACTN4 knockout cells used as negative cell line), MCF7 IP: HeLa cell lysate
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR2533(2)

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab108198于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (4)	1/1000 - 1/10000. Detects a band of approximately 100 kDa (predicted molecular weight: 105 kDa).
IP		1/30.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF	★★★★★ (3)	1/100 - 1/250.
Flow Cyt (Intra)		1/50. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

靶标

功能	F-actin cross-linking protein which is thought to anchor actin to a variety of intracellular structures. This is a bundling protein. Probably involved in vesicular trafficking via its association with the CART complex. The CART complex is necessary for efficient transferrin receptor recycling but not for EGFR degradation.
组织特异性	Widely expressed.
疾病相关	Defects in ACTN4 are the cause of focal segmental glomerulosclerosis type 1 (FSGS1) [MIM:603278]. A renal pathology defined by the presence of segmental sclerosis in glomeruli and resulting in proteinuria, reduced glomerular filtration rate and edema. Renal insufficiency often progresses to end-stage renal disease, a highly morbid state requiring either dialysis therapy or kidney transplantation.
序列相似性	Belongs to the alpha-actinin family. Contains 1 actin-binding domain. Contains 2 CH (calponin-homology) domains. Contains 2 EF-hand domains. Contains 4 spectrin repeats.
细胞定位	Nucleus. Cytoplasm. Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Colocalizes with actin stress fibers. Nuclear translocation can be induced by the PI3 kinase inhibitor wortmannin or by cytochalasin D. Exclusively localized in the nucleus in a limited number of cell lines.

图片



Western blot - Anti-alpha Actinin 4 antibody
[EPR2533(2)] (ab108198)


Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198) at 1/10000 dilution (Purified) + HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate at 15 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 105 kDa

Observed band size: 105 kDa



Western blot - Anti-alpha Actinin 4 antibody
[EPR2533(2)] (ab108198)

All lanes : Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198) at 1/1000 dilution (Purified)

Lane 1 : Human skeletal muscle lysate

Lane 2 : A431 (Human epidermoid carcinoma epithelial cell) whole cell lysate

Lane 3 : Mouse brain lysate

Lane 4 : Rat brain lysate

Lane 5 : Rat heart lysate


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution


Predicted band size: 105 kDa

Observed band size: 105 kDa




Flow Cytometry (Intracellular) - Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198)

Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labelling alpha Actinin 4 with Purified ab108198 at 1:50 dilution (10 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) secondary antibody was used at 1:2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabelled control - Cell without incubation with primary antibody and secondary antibody (Blue).




Immunocytochemistry/ Immunofluorescence - Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198)

Immunocytochemistry analysis of MCF7 (Human breast adenocarcinoma epithelial cell) cells labeling alpha Actinin 4 with Purified ab108198 at 1:250 dilution (2.0 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat liver tissue sections labeling alpha Actinin 4 with Purified ab108198 at 1:150 dilution (3.29 µg/mL). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver tissue sections labeling alpha Actinin 4 with Purified ab108198 at 1:150 dilution (3.29 µg/mL). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue sections labeling alpha Actinin 4 with Purified ab108198 at 1:150 dilution (3.29 µg/mL). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Immunocytochemistry/ Immunofluorescence - Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198)

ab108198 staining ACTN4 in wild-type HAP1 cells (top panel) and ACTN4 knockout HAP1 cells (bottom panel). The cells were fixed with PFA (10min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab108198 at 1/100 dilution and **ab7291** (Tubulin) at 1/1000 dilution overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (**ab150120**) at 2 µg/ml (shown in pseudo color red). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Western blot - Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198)

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)


Lane 2: ACTN4 (alpha Actinin 4) knockout HAP1 whole cell lysate (20 µg)

Lane 3: HeLa whole cell lysate (20 µg)

Lane 4: MCF7 whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab108198 observed at 105 kDa. Red - loading control, [ab9484](#), observed at 37 kDa.

ab108198 was shown to specifically react with alpha Actinin 4 in wild-type HAP1 cells as signal was lost in ACTN4 (alpha Actinin 4) knockout cells. Wild-type and ACTN4 (alpha Actinin 4) knockout samples were subjected to SDS-PAGE. ab108198 and [ab9484](#) (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Immunoprecipitation - Anti-alpha Actinin 4 antibody [EPR2533(2)] (ab108198)

Purified ab108198 at 1/50 dilution (2µg) immunoprecipitating alpha Actinin 4 in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10µg

Lane 2 (+): ab108198 + HeLa whole cell lysate.

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of ab108198 in HeLa whole cell lysate.

VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) (1/1000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm/TBST.

Observed band size: 105 kDa

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-alpha Actinin 4 antibody [EPR2533(2)]
(ab108198)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors