

Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free ab210714

敲除验证
重组
RabMAb

10 图像

概述

产品名称	Anti-AMPK alpha 1抗体[Y365] - BSA and Azide free
描述	兔单克隆抗体[Y365] to AMPK alpha 1 - BSA and Azide free
宿主	Rabbit
特异性	This antibody is specific for human AMPK alpha 1. This antibody shows low affinity on mouse and rat samples.
经测试应用	适用于: Flow Cyt (Intra), WB, IP, IHC-P, ICC/IF
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HeLa, HepG2, C6, NIH/3T3 and MCF-7 cell lysate. Mouse liver, brain, retina, and skeletal muscle tissue lysates. IHC-P: Human cervical carcinoma and lung carcinoma tissues. ICC/IF: MCF-7 cells. Flow Cyt (intra): HeLa cells. IP: HeLa cell lysate.
常规说明	ab210714 is the carrier-free version of ab32047 .

Our **carrier-free** antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our **conjugation kits** for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.20 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	Y365
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab210714于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		Use at an assay dependent concentration. Predicted molecular weight: 63 kDa.
IP		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols</u> .
ICC/IF		Use at an assay dependent concentration.

靶标

功能	Responsible for the regulation of fatty acid synthesis by phosphorylation of acetyl-CoA carboxylase. It also regulates cholesterol synthesis via phosphorylation and inactivation of hormone-sensitive lipase and hydroxymethylglutaryl-CoA reductase. Appears to act as a metabolic stress-sensing protein kinase switching off biosynthetic pathways when cellular ATP levels are depleted and when 5'-AMP rises in response to fuel limitation and/or hypoxia. This is a catalytic subunit.
序列相似性	Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. SNF1 subfamily. Contains 1 protein kinase domain.

图片



Western blot - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

All lanes : Anti-AMPK alpha 1 antibody [Y365] ([ab32047](#)) at 1/1000 dilution

Lane 1 : Wild-type RAW 264.7 cell lysate

Lane 2 : PRKAA1 knockout RAW 264.7 cell lysate

Lane 3 : Mouse Liver cell lysate


Lane 4 : Neuro2A cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 63 kDa


False colour image of Western blot: Anti-AMPK alpha 1 antibody [Y365] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab32047](#) was shown to bind specifically to AMPK alpha 1. A band was observed at 64 kDa in wild-type RAW 264.7 cell lysates with no signal observed at this size in PRKAA1 knockout cell line [ab280055](#) (knockout cell lysate [ab280114](#)). To generate this image, wild-type and PRKAA1 knockout RAW 264.7 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling AMPK alpha 1 with purified **ab32047** at 1/100. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).




Immunocytochemistry/ Immunofluorescence - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Immunocytochemistry/Immunofluorescence analysis of MCF7 cells labelling AMPK alpha 1 with purified **ab32047** at 1/250. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.

Control 1: primary antibody (1/250) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).



Flow Cytometry (Intracellular) - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Intracellular Flow Cytometry analysis of HeLa cells labelling AMPK alpha 1 with purified **ab32047** at 1/150 (red). Cells were fixed with 2% paraformaldehyde. A FITC-conjugated goat anti-rabbit IgG (1/150) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).



Immunoprecipitation - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

ab32047 (purified) at 1/40 immunoprecipitating AMPK alpha 1 in HeLa whole cell lysate.

Lane 1 (input): HeLa whole cell lysate (10µg)

Lane 2 (+): **ab32047** + HeLa whole cell lysate (10µg).


Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of **ab32047** in HeLa whole cell lysate.

For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/1500 dilution.

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).



Western blot - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

All lanes : Anti-AMPK alpha 1 antibody [Y365] ([ab32047](#)) at 1/500 dilution

Lane 1 : HepG2 (human liver hepatocellular carcinoma cell line) whole cell lysate

Lane 2 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate

Lane 3 : Mouse liver tissue lysate

Lane 4 : Mouse brain tissue lysate

Lane 5 : Mouse kidney tissue lysate

Lane 6 : Mouse retina tissue lysate

Lane 7 : Mouse skeletal muscle tissue lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 63 kDa


Observed band size: 63 kDa

Additional bands at: 40 kDa (possible non-specific binding)

Exposure time: 3 minutes

Blocking/Diluting buffer and concentration: 5% NFDM/TBST.

This antibody shows low affinity on mouse samples. This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab32047](#)).



All lanes : Anti-AMPK alpha 1 antibody [Y365] ([ab32047](#)) at 1/20000 dilution

Lane 1 : MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysate

Lane 2 : HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate

Lane 3 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

Lane 4 : K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate

Lane 5 : C6 (Rat glial tumor glial cell) whole cell lysate

Lane 6 : Neuro-2a (Mouse neuroblastoma neuroblast) whole cell lysate

Lane 7 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 63 kDa


Observed band size: 63 kDa

Exposure time: 180 seconds

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

This antibody shows low affinity on mouse and rat samples.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab32047](#)).



Flow Cytometry (Intracellular) - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Overlay histogram showing HeLa cells stained with unpurified **ab32047** (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (unpurified **ab32047**, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (**ab96899**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cervical carcinoma tissue labelling AMPK alpha 1 with unpurified **ab32047** at a dilution of 1/100.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32047**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-AMPK alpha 1 antibody [Y365] - BSA and Azide free (ab210714)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors