# abcam

### Product datasheet

# Anti-Annexin-4/ANXA4 antibody [EPR22929-208] ab256456



重组 RabMAb

1 References 16 图像

概述

产品名称 Anti-Annexin-4/ANXA4抗体[EPR22929-208]

描述 兔单克隆抗体[EPR22929-208] to Annexin-4/ANXA4

宿主 Rabbit

经测试应用 适用于: IHC-Fr, ICC/IF, IHC-P, WB, Flow Cyt (Intra)

种属反应性 与反应: Mouse, Rat, Human

免疫原 Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

阳性对照 WB: Human kidney, liver and lung lysate. HepG2, HeLa, C6, RAW 264.7, PC-12 and NIH/3T3

> lysates. Mouse liver, kidney and lung lysate. Rat kidney lysate. IHC-P: Human pancreas, kidney, endometrium, mouse and rat stomach tissue. ICC/IF: HepG2 and NIH/3T3 cells. Flow Cyt (intra):

HepG2 and NIH/3T3 cells.

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

存储溶液 pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)

纯度 Protein A purified

克隆 单克隆

**克隆编号** EPR22929-208

同种型 IgG

#### 应用

The Abpromise guarantee Abpromise™承诺保证使用ab256456于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-Fr		1/100. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).
ICC/IF		1/100.
IHC-P		1/5000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		1/2000. Predicted molecular weight: 35 kDa.
Flow Cyt (Intra)		1/500.

应用说明 Is unsuitable for IP.

靶标

功能 Calcium/phospholipid-binding protein which promotes membrane fusion and is involved in

exocytosis.

**序列相似性** Belongs to the annexin family.

Contains 4 annexin repeats.

结**构域** A pair of annexin repeats may form one binding site for calcium and phospholipid.

图片



[EPR22929-208] (ab256456)

All lanes: Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456) at 1/2000 dilution

Lane 1: C6 (rat glial tumor glial cell) whole cell lysate

Lane 2: RAW 264.7 (mouse abelson murine leukemia virus-

induced tumor macrophage) whole cell lysate

Lane 3: PC-12 (rat adrenal gland pheochromocytoma) whole cell

lysate

Lane 4: NIH/3T3 (mouse embryonic fibroblast) whole cell lysate

Lysates/proteins at 10 µg per lane.

#### Secondary


All lanes: Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at

1/100000 dilution

Predicted band size: 35 kDa Observed band size: 31,36 kDa

The expression molecular weight observed is consistent with what has been described in the literature (PMID: 16893879). Blocking/Diluting buffer and concentration: 5% NFDM/TBST.

Exposure Time: 15 seconds.



Lane 1: Human kidney lysate

Lane 2: Human liver lysate

Lane 3: Human lung lysate

Lane 4: HepG2 (human hepatocellular carcinoma epithelial cell)

whole cell lysate

Lane 5: HeLa (human cervix adenocarcinoma epithelial cell) whole

cell lysate

Lane 6: Mouse liver lysate


Lane 7: Mouse kidney lysate

Lane 8: Mouse lung lysate

Lane 9: Rat kidney lysate

Lysates/proteins at 20 µg per lane.

# Secondary



Western blot - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)

**Lanes 1-3 :** VeriBlot for IP Detection Reagent (HRP) (<u>ab131366</u>) at 1/1000 dilution


Lanes 4-9: Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

**Predicted band size:** 35 kDa **Observed band size:** 31,36 kDa

The expression molecular weight observed is consistent with what has been described in the literature (PMID: 16893879).

Blocking/Diluting buffer and concentration: 5% NFDM/TBST.

Exposure Time: Lanes 1-4: 59 seconds; Lanes 5-8: 125 seconds; Lane 9: 10 seconds.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)

Immunohistochemical analysis of paraffin-embedded rat stomach tissue labeling Annexin-4/ANXA4 with ab256456 at 1/5000 dilution (0.103 µg/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining in rat stomach. The section was incubated with ab256456 for 10 mins at RT. The immunostaining staining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)

Immunohistochemical analysis of paraffin-embedded mouse stomach tissue labeling Annexin-4/ANXA4 with ab256456 at 1/5000 dilution (0.103 µg/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining in mouse stomach. The section was incubated with ab256456 for 10 mins at RT. The immunostaining staining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)


Immunohistochemical analysis of paraffin-embedded human endometrium tissue labeling Annexin-4/ANXA4 with ab256456 at 1/5000 dilution (0.103 µg/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining in human endometrium (PMID: 16954445) is observed. The section was incubated with ab256456 for 10 mins at RT. The immunostaining staining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)



Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)


Immunohistochemical analysis of paraffin-embedded human pancreas tissue labeling Annexin-4/ANXA4 with ab256456 at 1/5000 dilution (0.103 µg/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining in centroacinar cells and islets of human pancreas. The section was incubated with ab256456 for 10 mins at RT. The immunostaining staining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.



Flow Cytometry (Intracellular) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456) Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labelling Annexin-4/ANXA4 with ab256456 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) / Black isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor<sup>®</sup> 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Flow Cytometry (Intracellular) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456) Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HepG2 (human hepatocellular carcinoma epithelial cell) cells labelling Annexin-4/ANXA4 with ab256456 at 1/500 dilution (Red) compared with a Rabbit monoclonal lgG (ab172730) / Black isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit lgG (Alexa Fluor<sup>®</sup> 488, ab150077) at 1/2000 dilution was used as the secondary antibody.



Immunohistochemistry (Frozen sections) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456) Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized rat pancreas tissue labeling Annexin-4/ANXA4 with ab256456 at 1/100 dilution, followed by **ab150077** AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 dilution (green). Positive staining on mouse pancreas (PMID: 25176043) is observed. The nuclear counter stain is DAPI (blue). Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab150077** AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 dilution.


Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Immunohistochemistry (Frozen sections) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)

Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized mouse pancreas tissue labeling Annexin-4/ANXA4 with ab256456 at 1/100 dilution, followed by **ab150077** AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 dilution (green). Positive staining on mouse pancreas (PMID: 25176043) is observed. The nuclear counter stain is DAPI (blue). Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab150077** AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 dilution.


Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Immunohistochemistry (Frozen sections) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)

Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized rat kidney tissue labeling Annexin-4/ANXA4 with ab256456 at 1/100 dilution, followed by **ab150077** AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 dilution (green). Positive staining on rat kidney (PMID: 15145526) is observed. The nuclear counter stain is DAPI (blue). Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab150077** AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 dilution.


Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Immunohistochemistry (Frozen sections) - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)

Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized mouse kidney tissue labeling Annexin-4/ANXA4 with ab256456 at 1/100 dilution, followed by **ab150077** AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 dilution (green). Positive staining on mouse kidney (PMID: 15145526) is observed. The nuclear counter stain is DAPI (blue). Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab150077** AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 dilution.


Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Immunocytochemistry/ Immunofluorescence - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labeling Annexin-4/ANXA4 with ab256456 at 1/100 5.2  $\mu$ g/ml dilution, followed by **ab150077** AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary antibody at 1/1000 2  $\mu$ g/ml dilution (Green). Confocal image showing nuclear, cytoplasmic and staining in NIH/3T3 cell line. **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor<sup>®</sup> 594) was used to counterstain tubulin at 1/200 dilution (Red). The nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is  $\underline{ab150077}$  AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 2  $\mu$ g/ml dilution.



Immunocytochemistry/ Immunofluorescence - Anti-Annexin-4/ANXA4 antibody [EPR22929-208] (ab256456) Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HepG2 (human hepatocellular carcinoma epithelial cell) cells labeling Annexin-4/ANXA4 with ab256456 at 1/100 5.2 µg/ml dilution, followed by ab150077 AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 2 µg/ml dilution (Green). Confocal image showing nuclear, cytoplasmic and staining in HepG2 cell line. ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is  $\underline{ab150077}$  AlexaFluor<sup>®</sup>488 Goat anti-Rabbit secondary at 1/1000 2  $\mu$ g/ml dilution.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

## Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors