abcam

Product datasheet

APC Anti-CD14 antibody [MEM-18] ab239289

1 图像

概述

产品名称 APC Anti-CD14抗体[MEM-18]

描述 APC小鼠单克隆抗体[MEM-18] to CD14

宿主 Mouse

偶联物 APC. Ex: 645nm, Em: 660nm

 经测试应用
 适用于: Flow Cyt

 种属反应性
 与反应: Human

预测可用于: Non human primates 4

免疫原 Full length native protein (purified) corresponding to Human CD14. (A crude mixture of human

urinary proteins precipitated by ammonium sulphate from the urine of a patient suffering from

proteinuria).

阳性对照 Flow Cyt: Human peripheral blood leukocytes.

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C. Store In the Dark.

存储溶液 pH: 7.4

Preservative: 0.0975% Sodium azide

Constituent: PBS

纯**度** Size exclusion

纯**化说明** Purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum

conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct

use. No reconstitution is necessary.

1

 克隆
 单克隆

 克隆编号
 MEM-18

 同种型
 IgG1

应用

The Abpromise guarantee Abpromise™承诺保证使用ab239289于以下的经测试应用


"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use 10µl for 10 ⁶ cells. (or 100 µl of whole blood).

靶标

功能	Cooperates with MD-2 and TLR4 to mediate the innate immune response to bacterial lipopolysaccharide (LPS). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Up-regulates cell surface molecules, including adhesion molecules.	
组织 特异性	Expressed strongly on the surface of monocytes and weakly on the surface of granulocytes; also expressed by most tissue macrophages.	
序列相似性	Contains 11 LRR (leucine-rich) repeats.	
翻译后修饰	N- and O- glycosylated. O-glycosylated with a core 1 or possibly core 8 glycan.	
细胞定位	Cell membrane.	

图片



Flow cytometric analysis of human peripheral blood leukocytes labeling CD14 with ab239289. Surface staining.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors