

APC Anti-CD45 antibody [MEM-28] ab28106

★★★★★ [1 Abreviews](#) [5 References](#) [3 图像](#)


概述

产品名称	APC Anti-CD45抗体[MEM-28]
描述	APC小鼠单克隆抗体[MEM-28] to CD45
宿主	Mouse
偶联物	APC. Ex: 645nm, Em: 660nm
特异性	ab28106 reacts with all alternative forms of human CD45 antigen
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Human
免疫原	Tissue, cells or virus corresponding to Human CD45. Human thymocytes and T lymphocytes
阳性对照	Flow Cyt: Human peripheral blood cells.
常规说明	<p>The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能


形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	<p>pH: 7.4</p> <p>Preservative: 0.097% Sodium azide</p> <p>Constituents: PBS, 0.2% BSA</p>
纯度	IgG fraction
克隆	单克隆
克隆编号	MEM-28

同种型	IgG1							
应用	<p>The Abpromise guarantee Abpromise™承诺保证使用ab28106于以下的经测试应用</p> <p>“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。</p> <table> <tr> <th>应用</th><th>Ab评论</th><th>说明</th></tr> <tr> <td>Flow Cyt</td><td>★★★★★ (1)</td><td>Use at an assay dependent concentration.</td></tr> </table>		应用	Ab评论	说明	Flow Cyt	★★★★★ (1)	Use at an assay dependent concentration.
应用	Ab评论	说明						
Flow Cyt	★★★★★ (1)	Use at an assay dependent concentration.						
靶标								
功能	<p>Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN.</p>							
疾病相关	<p>Defects in PTPRC are a cause of severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive (T(-)B(+)NK(+) SCID) [MIM:608971]. A form of severe combined immunodeficiency (SCID), a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity, leukopenia, and low or absent antibody levels. Patients present in infancy recurrent, persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development. Genetic variations in PTPRC are involved in multiple sclerosis susceptibility (MS) [MIM:126200]. MS is a neurodegenerative disorder characterized by the gradual accumulation of focal plaques of demyelination particularly in the periventricular areas of the brain. Peripheral nerves are not affected. Onset usually in third or fourth decade with intermittent progression over an extended period. The cause is still uncertain.</p>							
序列相似性	<p>Belongs to the protein-tyrosine phosphatase family. Receptor class 1/6 subfamily.</p> <p>Contains 2 fibronectin type-III domains.</p> <p>Contains 2 tyrosine-protein phosphatase domains.</p>							
结构域	<p>The first PTPase domain interacts with SKAP1.</p>							
翻译后修饰	<p>Heavily N- and O-glycosylated.</p>							
细胞定位	<p>Membrane. Membrane raft. Colocalized with DPP4 in membrane rafts.</p>							
图片								



Flow cytometry surface staining pattern of human peripheral whole blood stained using ab28106 at (10 μ l reagent / 100 μ l of peripheral whole blood).


Flow Cytometry - APC Anti-CD45 antibody [MEM-28] (ab28106)



ab28106 staining the CD45 in Human PBMCs by Flow Cytometry. Cells were prepared by Ficoll-Hypaque isolation of PBMCs from buffy coat. The sample was incubated with the primary antibody (1/25 in PBS + 2% Human serum and 1mM EDTA) for 25 minutes at 4°C.

Flow Cytometry - APC Anti-CD45 antibody [MEM-28] (ab28106)

This image is courtesy of an anonymous Abreview



Separation of human CD45 positive lymphocytes (red-filled) from human CD45 negative blood debris (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using ab28106 (10 µl reagent / 100 µl of peripheral whole blood).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors